

Isophotal shapes of early-type galaxies to very faint levels

Laxmikant Chaware,¹ Russell Cannon,² Ajit K. Kembhavi,³ Ashish Mahabal⁴

and

S. K. Pandey¹

ABSTRACT

We report on a study of the isophotal shapes of early-type galaxies, to very faint levels reaching $\sim 0.1\%$ of the sky brightness. The galaxies are from the Large Format Camera (LFC) fields obtained using the Palomar 5 m Hale telescope, with integrated exposures ranging from 1 to 4 hours in the SDSS r , i and z bands. The shapes of isophotes of early-type galaxies are important as they are correlated with the physical properties of the galaxies and are influenced by galaxy formation processes. In this paper we report on a sample of 132 E and SO galaxies in one LFC field. We have redshifts for 53 of these, obtained using AAOmega on the Anglo-Australian Telescope. The shapes of early-type galaxies often vary with radius. We derive average values of isophotal shape parameters in four different radial bins along the semi-major axis in each galaxy. We obtain empirical fitting formulae for the probability distribution of the isophotal parameters in each bin and investigate for possible correlations with other global properties of the galaxies. Our main finding is that the isophotal shapes of the inner regions are statistically different from those in the outer regions. This suggests that the outer and inner parts of early-type galaxies have evolved somewhat independently.

Subject headings: galaxies: elliptical and lenticular, cD — galaxies: photometry — galaxies: statistics — galaxies: structure

1. Introduction

The morphology and colours of galaxies carry important information about their formation and evolution. Bender et al. (1988) studied a sample of 109 large, nearby early-type

¹School of Studies in Physics, Pt. Ravishankar Shukla University, Raipur 492010, India

²Australian Astronomical Observatory, PO Box 915, North Ryde, NSW 1670, Australia.

³Inter University Center for Astronomy and Astrophysics, Pune, India

⁴California Institute of Technology, Pasadena, CA 91125, USA

galaxies to examine their isophotal shapes, using Fourier expansions in the polar angle to distinguish between boxy (rectangular) isophotes and disk (pointed) isophotes. Studies have shown that elliptical galaxies with disk isophotes tend to be fainter, rotationally supported, lack X-ray and radio activity and have power-law nuclear light profiles, while those with boxy isophotes tend to be brighter, supported by random motions, have significant X-ray and radio activity and peaked nuclear profile (Ferrarese et al. 1994; Bosch 1994; Rest et al. 2001; Lauer 2005). The shapes of galaxies may also be correlated with their ages (Ryden et al. 2001). Recently Hao et al. (2006) studied the isophotal shapes of a large sample of 847 early-type galaxies from the SDSS, and Pasquali et al. (2007) have used the same sample to investigate the dependence of the isophotal structure on the AGN activity and environment of the galaxies. The dynamic range available to these studies is limited, due to the short exposure of the SDSS. The isophotal shape study by Bender et al. (1988) extends to 1.5 times effective radius ($1.5r_e$) of the galaxies, while Hao et al. (2006) studied their larger sample of 857 nearby galaxies with redshift $z < 0.05$ extending to 1.5 times the Petrosian half-light radius ($1.5R_{50}$).

In this paper we revisit the properties of isophotal shapes for a sample of more distant early-type galaxies in a single field, using exceptionally deep CCD exposures. These enable us to go out to much larger radii of $\sim 4.5R_{50}$ and reach $\sim 4 \text{ mag arcsec}^{-2}$ deeper in surface brightness. We have obtained redshifts for a subsample of these galaxies: the redshifts peak at $z \sim 0.1$ and extend to $z \sim 0.8$, covering a wide range of absolute magnitudes.

Studies of the faint outer parts of galaxies are not new. By 1980 Malin (e.g. Malin & Carter 1980) was using sky survey photographs to create images that reached 10 magnitudes below the level of the sky background. These led to the discovery of very faint shells and tails around galaxies, soon interpreted as evidence for dynamical evolution due to gravitational interactions and mergers (e.g. Quinn 1984). Capaccioli et al. (1988) combined photographic and CCD imaging of nine nearby galaxies, to explore the boundary between E and S0 galaxies by looking at the changing shapes of isophotes with radius.

We extend the methodology of Bender et al. (1988) and derive mean isophote parameters for up to four radial bins for each galaxy. Pointed isophotes can usually be attributed to a galaxy which consists of a spheroid and a weak disk which is being viewed more or less edge-on. Nieto & Bender (1989) suggest that some pointed isophotes may be due to tidal extensions. But the origin of boxy isophotes is still unclear. Naab et al. (1999) showed that mergers of spiral galaxies with comparable mass are more likely to form boxy ellipticals, while mergers of spiral galaxies with differing mass can lead to disk ellipticals. However, disc-disc mergers do not form a perfect dichotomy (Nabb & Burkert 2003), while elliptical-elliptical mergers always lead to boxy isophotes (Nabb & Burkert 2006). Using N-body simulation

Bournaud et al. (2005) found that for a 7:1 merger, isophotes in the inner region (bulge) are boxy, while the outer isophotes are disk. Such radial variation is also seen in the analysis of 2MASS data for Arp mergers by Chitre & Jog (2002). Our study of isophotal shapes to faint levels will be useful for better comparison between observations and such theoretical possibilities.

This paper is organized in the following way: in Section 2 we describe the imaging data, sample selection, bulge-disk decomposition and spectroscopic data used for this study. In Section 3, surface photometry of sample galaxies, and estimation of isophotal and rest frame parameters, are described. In Section 4, we give the properties of the sample and main results of our study along with the comparison with earlier work. We discuss our results in Section 5 and give a summary and our conclusions in Section 6.

2. The observational data

2.1. Imaging data

The imaging data were originally obtained as part of a faint quasar survey using the Large Format Camera (LFC) on the 5 m Hale telescope at Mount Palomar Observatory, with integrated exposures ranging from 1 to 4 hours in r , i and z filters. The LFC is a mosaic of six 2048×4096 pixel CCDs, mounted at the prime focus of the Hale Telescope. The fields targeted were those of known SDSS high redshift ($z > 5$) quasars in order to look for a local over-density of quasars (Mahabal et al. 2005). We obtained reduced images which were bias subtracted, flat fielded and processed to remove cosmic rays as a part of the original programme. The deep LFC images have excellent signal-to-noise ratio (SNR), permitting us to examine faint features, including the outermost extensions of the galaxies, down to $\sim 0.1\%$ of the sky brightness.

In this paper we use a sample of galaxies selected from the LFC field SDSS 1208+0010 which spans an area of $13'.56 \times 25'.32$ centered on $RA = 12^h08^m24^s$ and $dec = 00^d10^m28^s$ (J2000). The field was observed in the SDSS i and z filters with integrated exposure of 2.51 h and 2.56 h respectively. The i -band image of the field is an average combination of 140 individual frames. The mean background count in the i -band image is 6278 ADU (analog-to-digital units), so with a gain of 1.1, an accuracy of $\sim 0.1\%$ is obtained in the estimation of the sky background.

All the LFC fields are in the area of the sky covered by SDSS imaging. We therefore have imaging data in the SDSS u , g and r bands for our sample galaxies, in addition to the LFC i and z band data, though not to the same depth. We have used the SDSS DR7

photometric catalog (Abazajian et al. 2009) to obtain the Petrosian half-light radii, apparent magnitudes and Galactic extinction corrections at the position of each sample galaxy in all the five bands.

2.2. Sample selection

We used SExtractor (Bertin & Arnouts 1996) to produce a catalog of all sources in the LFC field SDSS 1208+0010, and to identify galaxies from the catalog. SExtractor provides a *stellarity index* for star/galaxy separation of the detected objects: 0 for an object identified as a galaxy, 1 for a star, and some intermediate value for ambiguous objects, which are usually the fainter objects in the sample. For correct star/galaxy separation, and to have galaxies with sufficient brightness and size to ensure reliable morphological study, we select galaxies using objects from our SExtractor catalog which satisfy the following criteria: (i) SExtractor parameter value CLASS_STAR < 0.8; (ii) SDSS parameter value PhotoType = GALAXY in the PhotoObjAll table of the SDSS DR7 photometric catalog (Abazajian et al. 2009); and (iii) $m_i < 20.5$ where m_i is the i -band model magnitude of the object corrected for extinction, which is denoted by *dered_i* in the SDSS DR7 photometric catalog. To ensure that a sufficient number of pixels is available for surface photometry, we included in our sample only objects with the i -band SExtractor parameter ISOAREA_IMAGE > 165. The area covered by the chosen pixels is > 21.74 arcsec² for every object in the sample, with the intensity in every pixel being brighter than the analysis threshold of 26.0 mag arcsec⁻², which corresponds to a 3 σ detection threshold. Through visual examination of the image, we have excluded objects which are near bright stars. Applying these criteria, we arrived at a sample of 266 galaxies.

Basic data for the target galaxies in Field SDSS 1208+0010 are available in Table 9, which is at the end of this article. We give first few entries from Table 9 in Table 1 as an illustration. The table gives an LFC serial number for each galaxy, from the SExtractor catalog generated by us, the SDSS ID, position and photometric data from the DR7 photometric catalog (Abazajian et al. 2009), and finally the semi-major axis (in arcsec), i -band surface brightness (in mag arcsec⁻²) and signal-to-noise ratio (SNR) of the outermost isophote to which an ellipse can be fitted (see Section 3).

We show in Figure 1 the i -band image of the largest galaxy from the LFC field, together with the SDSS i -band image of the same galaxy. The profiles of the surface brightness along the semi-major axis of the two images are shown in the lower panels, up to a point where the ratio of the mean isophotal intensity to the total error in isophotal intensity is about three. These points are shown as arrows along the two profiles, with the surface brightness level reached at that point given as a fraction of the sky brightness. Since the SNR is the same for




Fig. 1.— Surface brightness (SB) profiles of a galaxy from the sample, generated using SDSS (shown in blue color) and LFC (shown in red color) i band images. The top panels show SDSS (left) and LFC (right) image cutouts ($\sim 55'' \times 55''$). The SB profiles are obtained by fitting ellipses up to a point along the semi major axis where the ratio of the mean isophotal intensity to the total error in the isophotal intensity drops to three in each image. 1σ error in the isophotal intensity is indicated for each point. The dotted vertical line shows the average seeing in the i band LFC observation. The galaxy has $m_i = 17.8$, and $z = 0.1772$ as determined from AAT observation. The linear scale for the cosmology adopted (see Section 3) is $\sim 3 \text{ kpc arcsec}^{-1}$.

isophotes of the two images at the indicated points, it is clear that we reach $\sim 4 \text{ mag arcsec}^{-2}$ fainter in surface brightness in the LFC images compared to the SDSS images (see Section A). The mean value of the semi-major axis (sma) length for the outermost isophote for our sample galaxies is $5''.41$. The ellipse fitting process fails to converge before reaching the condition that isophotes have (mean isophotal intensity/total error in isophotal intensity) ~ 3 for most of the cases, principally because of crowding effects at the lowest brightness levels. For the sample the mean signal-to-noise ratio reached at the outermost isophote is 14. The average surface brightness and its average error at the outermost isophote we could reach for our sample galaxies are $26.94 \text{ mag arcsec}^{-2}$ and $0.12 \text{ mag arcsec}^{-2}$ respectively.

2.3. Bulge-disk decomposition

We have performed bulge-disk decomposition and selected 132 galaxies with bulge-to-total luminosity ratio $B/T > 0.4$ as being of early-type (see Rawat et al. 2007, for a discussion), which will form the basis for work reported in this paper. Here the luminosities are estimated using parameters which define the large scale structure of the galaxy. We assume that each galaxy consists of a bulge and a disk, and used the two-dimensional code GALFIT (Peng et al. 2002) to decompose the image of the galaxy into these components. The intensity profile of the bulge is modeled with the Sersic law (Sersic 1968),

$$\Sigma(r) = \Sigma_e \exp \left[-2.303 b_n \left[\left(\frac{r}{r_e} \right)^{1/n} - 1 \right] \right], \quad (1)$$

where r_e is the half light radius, Σ_e the surface brightness at r_e and n the Sersic index. The disk is modeled with an exponential profile (Freeman 1970))

$$\Sigma(r) = \Sigma_0 \exp \left[-\frac{r}{r_d} \right], \quad (2)$$

where Σ_0 is the central surface brightness and r_d is the disk scale length. GALFIT obtains best fit parameters by minimizing χ^2 obtained by comparing the observed distribution of intensity in the two-dimensional image of the galaxy with model images obtained using the radial profiles defined above, and convolved with the point spread function. After the decomposition, the total bulge and disk fluxes are obtained by integrating the best fit profiles over all values of the semi-major axis.

2.4. Spectroscopic data

The distances to most of our galaxies are unknown. However, we have obtained spectra for some of the galaxies in the LFC 1208+0010 field using the 2dF/AAOmega multi-fiber

Table 1. Basic parametrs of the sample of early-type galaxies

	SDSS ID	RA	Dec	u mag	g mag	r mag	i mag	z mag	R_{50} arcsec	sma arcsec	μ_i mag arcsec ⁻²	SNR	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	
LFC 462	588848899913548358	182.02545140	-0.00014042	21.50 ± 0.43	21.06 ± 0.13	20.55 ± 0.13	20.57 ± 0.21	19.96 ± 0.43	1.07 ± 0.16	3.81	26.36 ± 0.06	17.25	
LFC 525	587748929241612656	182.10068456	0.00579127	22.48 ± 1.58	20.76 ± 0.09	20.42 ± 0.10	20.13 ± 0.12	20.74 ± 0.95	1.10 ± 0.11	3.81	26.06 ± 0.04	26.07	
LFC 554	587748929241547219	181.99182097	0.00889758	22.09 ± 1.04	21.89 ± 0.24	21.11 ± 0.18	19.90 ± 0.09	19.67 ± 0.34	1.28 ± 0.07	3.81	26.67 ± 0.08	12.50	
LFC 558	587748929241547253	181.99959155	0.01032991	26.27 ± 2.14	21.73 ± 0.24	20.20 ± 0.09	19.44 ± 0.07	19.05 ± 0.21	1.01 ± 0.07	5.11	26.58 ± 0.10	10.06	
LFC 622	587748929241546923	182.03581581	0.01761600	20.79 ± 0.41	19.26 ± 0.03	18.16 ± 0.02	17.70 ± 0.02	17.33 ± 0.06	1.38 ± 0.02	10.11	27.21 ± 0.09	11.88	
LFC 633	587748929241547046	182.04195384	0.01802068	20.48 ± 0.30	20.09 ± 0.06	19.51 ± 0.05	19.11 ± 0.05	19.21 ± 0.28	1.26 ± 0.05	8.32	27.18 ± 0.08	13.77	
LFC 663	587748929241612845	182.17111251	0.01778466	21.46 ± 0.60	21.97 ± 0.26	20.89 ± 0.15	20.06 ± 0.10	19.19 ± 0.22	1.14 ± 0.08	3.81	25.90 ± 0.05	23.48	
LFC 677	587748929241547621	181.99102394	0.02035316	19.94 ± 0.20	22.14 ± 0.42	20.94 ± 0.16	20.98 ± 0.33	22.79 ± 4.34	0.80 ± 0.17	5.36	27.64 ± 0.19	5.11	
LFC 725	587722983351583067	182.01578218	0.02274361	25.14 ± 4.14	21.58 ± 0.27	20.82 ± 0.13	20.75 ± 0.30	20.12 ± 0.77	1.32 ± 0.37	3.81	26.65 ± 0.12	8.92	
LFC 741	587748929241547216	181.99078515	0.02434438	21.46 ± 0.57	21.65 ± 0.19	20.74 ± 0.12	20.19 ± 0.12	19.80 ± 0.36	0.97 ± 0.09	4.86	27.60 ± 0.21	4.73	
LFC 824	587748929241547430	182.05600812	0.03170720	23.94 ± 10.03	23.14 ± 2.52	20.28 ± 0.18	20.44 ± 0.65	19.60 ± 0.78	0.74 ± 0.33	3.81	25.93 ± 0.04	28.48	
LFC 848	587748929241547282	182.00945210	0.03522617	22.05 ± 1.24	20.93 ± 0.14	20.26 ± 0.11	20.13 ± 0.17	19.63 ± 0.39	1.00 ± 0.12	3.81	26.89 ± 0.09	11.38	
LFC 925	587748929241612771	182.14531708	0.03984272	21.09 ± 0.52	21.00 ± 0.13	20.09 ± 0.07	19.76 ± 0.09	19.08 ± 0.24	1.27 ± 0.07	3.81	26.10 ± 0.05	21.72	
LFC 954	587748929241612750	182.13801353	0.04149656	24.88 ± 5.75	23.04 ± 0.87	20.93 ± 0.17	20.11 ± 0.14	20.28 ± 0.78	1.06 ± 0.15	3.63	26.12 ± 0.08	12.67	↗
LFC 985	587748929241547693	182.03813760	0.04718642	25.96 ± 2.24	25.08 ± 1.99	20.94 ± 0.14	19.70 ± 0.08	19.39 ± 0.25	1.21 ± 0.10	6.21	28.07 ± 0.20	4.93	

Note. — Here we list data for only the first 15 galaxies in our sample: the full table is available at the end of the article as Table 9. Column (1) gives ID of the sample galaxies in SExtractor catalog generated by us. Quantities from column (2) to column (10) are obtained from the PhotoObjAll table of the SDSS DR7 photometric catalog (Abazajian et al. 2009). Column (2) gives a unique SDSS identifier which is listed as objID in the PhotoObjAll table Columns (3) and (4) give J2000 right ascension and declination in degree. Columns (5) to (9) give extinction corrected Petrosian magnitudes in u , g , r , i and z bands. Column (10) gives Petrosian half-light radius in a arcsec. Columns (11) to (13) give length of semi-major axis (in arcsec), i -band surface brightness (in $\text{mag}/\text{arcsec}^{-2}$) and signal to noise ratio respectively of the outermost isophote in which ellipse can be fitted (see Section 4).

system on the AAT. For these galaxies we can derive distances, absolute magnitudes and linear sizes, to facilitate comparison with other samples. These single-fiber spectra do not sample our galaxies consistently and were only intended for determining redshifts, but they do confirm that most of our early-type galaxies selected on the basis of B/T are indeed early-type galaxies.

We chose to observe the 266 selected galaxies from the field SDSS 1208+0010 (see Section 2.2) using the AAOmega multi-fiber spectrograph¹ (Saunders et al. 2004) on the 4 m Anglo-Australian Telescope (AAT). AAOmega, which has 392 fibers each with 2'' angular diameter, was used with gratings 385R and 580V in multi-object mode. The detectors used were EEV 4482 2k × 4k CCDs having pixel size 15 μm, allowing 10 pixels per fiber. The wavelength range covered by grating 580V is 3700 – 5700 Å with dispersion 1.0 Å pixel⁻¹, and by grating 385R is 5600 – 8800 Å with dispersion 1.6 Å pixel⁻¹. This setup of AAOmega provides a resolution of $R \sim 1300$.

The compactness of our field as compared to the 2 degree field of view of AAOmega, together with a 30'' restriction on the minimum separation between fibers, meant that it was not possible to observe all the 266 galaxies simultaneously. We used the latest version of the AAOmega specific *CONFIGURE* program² for automatic fiber allocation. This program uses a “simulated annealing” algorithm (Miszalski et al. 2006) to maximize the number of objects observed. We assigned priorities to our targets as a function of fiber magnitude in the *g* filter (*fiberMag_g*). We selected the positions of blank sky regions and suitable guide stars from the SDSS *PhotoObjAll* table, lying within the 2dF field but outside the central LFC field. SDSS images were visually inspected to ensure there was no contamination by other objects.

The publicly available *2DFDR*³ (Bailey et al. 2004) pipeline reduction system was used on each exposure to subtract bias and dark current and to carry out flat-fielding, sky-subtraction and wavelength-calibration. The blue and red-arm spectra were spliced together after the spectra in each exposure set were co-added. Redshifts were determined using a version of the *ZCODE* software in manual mode (see Cannon et al. 2006, sec 5.2 & 5.3). Each spectrum was visually inspected and assigned a redshift quality flag *Q*, having value from 0 to 6 based on (i) emission-line/absorption-line features identified in the spectrum and (ii) cross-correlation value estimated by the software. The cross-correlation was done with that template spectrum which matched best with each observed spectrum. A larger value of

¹See: <http://www.aao.gov.au/local/www/aaomega/>

²See: http://www.aao.gov.au/AAO/2df/aaomega/aaomega_software.html

³<http://www.aao.gov.au/AAO/2df/software.html#2dfdr>

Q signifies a more reliable redshift measurement. Redshift measurements with quality $Q \geq 3$ were considered reliable, while $Q < 3$ was assigned to unreliable redshifts or when no redshift was obtained. From experience with much larger samples of galaxies for other projects (e.g. Cannon et al. 2006), we believe that around 95% of redshifts flagged as reliable should be correct.

The median Petrosian half-light radius of our sample of galaxies is about $1.05''$, only slightly larger than the radius of the 2dF/AAOmega fibers. Thus for large nearby galaxies our spectra sample only the nucleus, while for small distant galaxies we integrate most of the light. The majority of our spectra were best fitted by an early-type galaxy template, confirming the validity of our selection criteria based on B/T . We defer further discussion of the spectra to a subsequent paper: here we simply use the spectra to determine redshifts and hence distances.

AAOmega observations for galaxies in field SDSS 1208 were carried out in service mode with exposures of 1 h (3×1200 sec) in June 2006 and for 1.5 h (3×1800 sec) in March 2007. In the observations of June 2006 only 135 galaxies could be targeted. We attained a mean S/N per pixel of ~ 4 in the blue and ~ 10 in the red spectra for targets with $fiberMag_g \sim 21$. We obtained reliable redshifts for 55/135 (40%) of the galaxies targeted and achieved 100% success for the brightest targets, i.e. for all 37 galaxies with $fiberMag_g < 21.2$. However, the success rate fell rapidly for fainter galaxies. We observed the field SDSS 1208 again in March 2007 with increased exposure time and could target 121 galaxies from the field, including all those not targetted in 2006 plus 58 that failed to yield a redshift in the first run. By combining the spectra from the two observing runs we determined reliable redshifts for an additional 17 galaxies. It should be noted that our success in determining redshifts is higher when emission lines are present, and that our fiber spectra sample only a $2''$ diameter patch of sky which may contain a substantial fraction of the light from small distant galaxies but only the core of nearby galaxies. Thus they do not give an unbiased sampling of the full set of galaxies or of their spectral types.

We show in Figure 2 the redshift distribution for all the galaxies in field SDSS 1208 for which we obtained redshifts. The redshifts of galaxies which we selected as early-type on the basis of B/T are shown by the shaded area. It is evident that the proportion of early-type galaxies increases with redshift, with late-type galaxies being more common for $z < 0.2$ while early-type galaxies are more common for $z > 0.2$.

We have obtained the rest frame absolute magnitudes in the u , g , r , i and z bands for the galaxies for which we have redshift measurements using the equation

$$M = m - DM(z) - k(z) - A, \quad (3)$$

where z is the redshift, m and M are apparent and absolute magnitudes respectively in each band and $k(z)$ and A are the k-correction and Galactic extinction in that band. The distance modulus $DM(z)$ is obtained using a flat Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ and $H_0 = 70 \text{ km sec}^{-1} \text{ Mpc}^{-1}$. The k-corrections have been obtained using the publicly available KCORRECT code (Blanton & Roweis 2007), and the extinction values are obtained from the DR7 photometric catalog which uses Galactic extinction maps published by Schlegel et al. (1998).

To compare our results with earlier work, we have obtained Johnson B and V magnitudes using the transformations (Smith et al. 2002)

$$\begin{aligned} B &= g + 0.47(g - r) + 0.17 \\ V &= g - 0.55(g - r) - 0.03. \end{aligned} \quad (4)$$

The apparent B , V magnitudes are converted to absolute magnitudes using Equation 3 and k-corrections in B and V bands are obtained from Poggianti (1997).

To summarize, from the two sets of AAT observations we obtained reliable redshifts for 129 of the 266 galaxies in our original photometric sample: of these, 53 galaxies are included in the early-type sample discussed in this paper. The redshift-dependent parameters are listed in the Table 2. In order to facilitate comparison with other samples of galaxies, plots of absolute B magnitude and Petrosian R_{50} radius against redshift are shown in Figure 3. The bulk of the galaxies have redshifts in the range $0.1 < z < 0.3$, with M_B between -17 and -21 and Petrosian R_{50} from two to six kpc. There are half a dozen dwarfs with $z < 0.1$ and a similar number of giants, mostly at high redshifts. The upper panel of Figure 3 shows a good correlation between Petrosian R_{50} and z for most of the galaxies.




Fig. 2.— Redshift distribution of the 129 galaxies from field sdss1208. The shaded area shows the distribution of 53 early type galaxies which are part of this work.




Fig. 3.— The lower panel show plot of redshift versus M_B for the 53 galaxies of our sample. See text for details. The lowest redshift galaxies are mostly dwarfs, while those in the high redshift tail are the brightest. The plot of redshift versus Petrosian R_{50} radius in the upper panel shows a good correlation. The dwarf galaxies (with $M_B > -17$) are indicated by large circles, and the most luminous ($M_B < -21$) with triangle symbols.

Table 2. Distance-dependent parameters for 53 galaxies with redshifts

ID	z	R_{50}	sma	M_u	M_g	M_r	M_i	M_z	M_B	M_V
		kpc	kpc	mag	mag	mag	mag	mag	mag	mag
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
LFC 558	0.4382	5.72 ± 0.37	28.97	-18.48	-21.90	-22.49	-22.85	-23.07	-21.45	-22.26
LFC 622	0.1580	3.77 ± 0.06	27.59	-19.24	-20.66	-21.43	-21.78	-22.11	-20.13	-21.12
LFC 633	0.1162	2.65 ± 0.11	17.50	-18.26	-18.73	-19.27	-19.54	-19.47	-18.31	-19.06
LFC 663	0.6136	7.72 ± 0.57	25.75	-21.39	-22.18	-23.40	-23.79	-24.52	-21.44	-22.88
LFC 824	0.2230	2.66 ± 1.20	13.68	-17.42	-17.94	-20.20	-19.88	-20.67	-16.71	-19.21
LFC 848	0.2255	3.62 ± 0.45	13.80	-18.46	-19.60	-20.01	-20.02	-20.48	-19.24	-19.86
LFC 985	0.4361	6.85 ± 0.54	35.11	-19.53	-19.47	-22.37	-23.17	-23.21	-17.93	-21.10
LFC 988	0.2788	3.32 ± 0.37	16.13	-17.20	-19.09	-19.92	-20.69	-21.08	-18.54	-19.58
LFC 1039	0.3632	4.69 ± 0.84	29.91	-19.70	-20.37	-20.85	-20.52	-18.35	-19.98	-20.66
LFC 1056	0.0636	1.47 ± 0.06	9.70	-16.05	-17.47	-18.02	-18.18	-18.19	-17.05	-17.81
LFC 1091	0.4359	5.78 ± 0.93	28.88	-18.56	-20.78	-22.00	-22.42	-22.94	-20.04	-21.48
LFC 1411	0.1588	2.80 ± 0.24	10.45	-17.72	-18.35	-19.06	-19.16	-19.15	-17.85	-18.77
LFC 1494	0.2919	5.61 ± 0.17	36.36	-20.60	-21.36	-22.13	-22.61	-22.69	-20.83	-21.81
LFC 1504	0.2716	4.98 ± 0.34	27.07	-18.55	-19.75	-20.50	-21.05	-20.73	-19.23	-20.19
LFC 1551	0.1829	2.90 ± 0.56	11.72	-19.07	-18.93	-19.85	-20.02	-21.28	-18.33	-19.47
LFC 1613	0.1161	2.37 ± 0.53	11.27	-17.10	-17.47	-17.99	-17.76	-17.65	-17.06	-17.79
LFC 1739	0.0332	0.57 ± 0.06	3.38	-9.69	-13.95	-14.83	-15.53	-16.20	-13.36	-14.46
LFC 1772	0.1775	3.37 ± 0.19	16.90	-16.78	-19.34	-20.00	-20.00	-20.20	-18.86	-19.73
LFC 1802	0.2925	4.73 ± 0.20	25.88	-19.77	-20.89	-21.30	-21.50	-21.32	-20.53	-21.15
LFC 2159	0.2387	3.72 ± 0.42	14.40	-18.33	-19.53	-20.60	-20.95	-22.08	-18.86	-20.15
LFC 2168	0.2480	3.88 ± 0.18	14.82	-19.68	-20.59	-20.94	-20.76	-20.94	-20.25	-20.81
LFC 2177	0.2047	3.62 ± 0.34	17.16	-18.26	-19.50	-20.12	-19.98	-20.39	-19.05	-19.87
LFC 2180	0.1024	1.11 ± 0.11	6.77	-16.46	-16.19	-16.87	-17.71	-18.57	-15.70	-16.59
LFC 2181	0.2479	3.36 ± 0.09	19.84	-19.46	-20.74	-21.28	-21.46	-21.78	-20.32	-21.07
LFC 2184	0.0357	0.93 ± 0.01	7.92	-18.01	-18.64	-18.91	-18.97	-19.04	-18.35	-18.82
LFC 2310	0.1931	3.07 ± 0.33	12.24	-19.22	-19.47	-20.02	-20.10	-20.61	-19.05	-19.80
LFC 2356	0.1193	4.06 ± 0.53	8.20	-16.73	-18.07	-18.40	-18.45	-18.74	-17.75	-18.28
LFC 2362	0.0592	1.49 ± 0.23	7.84	-15.51	-14.99	-16.61	-17.28	-17.74	-14.06	-15.91
LFC 2405	0.0899	1.40 ± 0.38	5.80	-12.28	-15.57	-16.73	-16.86	-17.96	-14.86	-16.24
LFC 2414	0.2869	5.81 ± 0.15	37.72	-21.21	-22.22	-22.97	-23.28	-23.56	-21.69	-22.66
LFC 2484	0.1887	3.55 ± 0.46	13.91	-19.48	-18.08	-19.31	-19.39	-19.11	-17.34	-18.79
LFC 2525	0.2051	4.00 ± 0.30	12.83	-18.26	-19.44	-19.90	-20.07	-20.00	-19.06	-19.72
LFC 2549	0.1264	2.97 ± 0.04	21.78	-19.02	-20.26	-20.93	-21.24	-21.49	-19.78	-20.66
LFC 2692	0.1295	2.31 ± 0.38	11.79	-14.39	-17.83	-17.90	-17.70	-19.00	-17.63	-17.90
LFC 2708	0.0184	0.28 ± 0.02	1.42	-8.13	-12.93	-14.19	-14.57	-12.69	-12.17	-13.65

3. Isophotal analysis and estimation of parameters

The early-type galaxies selected for our study are bulge dominated and therefore have isophotes which are close to elliptical in shape. It is our aim to measure morphological parameters which describe the elliptical shape, as well as small deviations from it, as a function of galaxy properties. We performed ellipse fitting using the IRAF routine `ellipse` within STSDAS, which is based on a technique described by Jedrzejewski (1987). However, before the ellipse fitting, pre-processing using SExtractor has to be performed to obtain a catalog of all detected objects in the field, a sky background image, and a *segmentation* image.

The segmentation image is a map of all detected objects in the field in which, for any given object, all the pixels have value equal to the running SExtractor catalog number corresponding to that object. The value of parameters used for estimation of background in the SExtractor configuration file are BACK_SIZE 128; BACK_FILTERSIZE 3; BACKPHOTO_TYPE LOCAL; and BACKPHOTO_THICK 64 (see Bertin & Arnouts 1996). The sky background image obtained is subtracted from the field image, and 150×150 pixel cutouts around the geometric center of our sample galaxies are made from the sky-subtracted image and the segmentation image. The cut-outs of segmentation images are used to generate mask files for the surface photometry.

We use the geometric center, ellipticity and position angle of sample galaxies, obtained from the SExtractor catalog, as initial values in the ellipse fitting. In the `ellipse` task, the image intensity is first sampled along a trial ellipse generated using these parameters, and the intensity string $I(\theta)$ is expanded in a Fourier series,

$$I(\theta) = I_0 + \sum_{n=1}^N [A_n \sin(n\theta) + B_n \cos(n\theta)], \quad (5)$$

where N is the highest harmonic fitted and θ is the azimuthal angle measured from the major axis. `ellipse` uses the first and second order coefficients obtained for trial ellipses to iteratively improve the fitting. For perfectly elliptical isophotes all the Fourier coefficients except I_0 , which is the mean isophotal intensity, should vanish at the end of the fitting process, and significant residuals can only be of order three or higher. We allowed the geometric center, ellipticity and position angle to vary freely during fitting. Successive ellipses are fitted along the semi-major axis with a logarithmic step of 0.05 until the ratio (mean isophotal intensity/total error in isophotal intensity) ~ 3 , or the ellipse fitting process fails to converge. The output of `ellipse` is a table containing radial profiles of number of isophotal parameters along with uncertainties associated with them (see Busko 1996, for error estimation of isophotal parameters).

Table 2—Continued

ID	z	R_{50}	sma	M_u	M_g	M_r	M_i	M_z	M_B	M_V
		kpc	kpc	mag	mag	mag	mag	mag	mag	mag
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
LFC 2712	0.1198	3.01 ± 0.17	16.30	-17.70	-19.21	-20.21	-20.74	-21.25	-18.58	-19.79
LFC 2782	0.1772	6.03 ± 0.13	40.61	-19.58	-20.83	-21.52	-21.87	-22.26	-20.33	-21.24
LFC 3072	0.4621	5.69 ± 0.57	29.86	-20.85	-20.65	-21.73	-21.94	-22.48	-19.97	-21.28
LFC 3094	0.2864	5.35 ± 0.21	32.55	-20.33	-21.19	-21.68	-21.80	-22.06	-20.79	-21.49
LFC 3218	0.2218	2.41 ± 0.16	16.56	-18.37	-19.02	-19.53	-19.74	-19.92	-18.61	-19.33
LFC 3274	0.0977	2.46 ± 0.31	6.56	-16.43	-16.74	-17.31	-17.69	-17.57	-16.30	-17.08
LFC 3292	0.4390	7.06 ± 0.56	35.24	-20.69	-21.76	-21.85	-22.06	-21.94	-21.54	-21.84
LFC 3381	0.7809	7.27 ± 0.55	38.03	-22.38	-22.79	-23.63	-23.38	-23.31	-22.23	-23.28
LFC 3430	0.1200	2.65 ± 0.10	24.12	-18.87	-19.89	-20.58	-20.91	-21.26	-19.40	-20.30
LFC 3458	0.3757	4.49 ± 0.32	25.14	-20.25	-20.69	-21.65	-22.01	-22.50	-20.07	-21.25
LFC 3508	0.2609	5.14 ± 0.23	35.25	-19.75	-20.97	-21.82	-22.27	-22.49	-20.41	-21.47
LFC 4009	0.1981	3.16 ± 0.21	20.33	-19.19	-20.31	-20.84	-21.12	-21.49	-19.89	-20.63
LFC 4059	0.3735	7.82 ± 0.40	52.06	-19.50	-21.67	-22.17	-22.43	-22.36	-21.27	-21.98
LFC 4188	0.0975	1.98 ± 0.05	10.66	-17.79	-18.99	-19.69	-19.97	-20.15	-18.50	-19.41
LFC 4248	0.0972	2.81 ± 0.19	14.96	-17.78	-18.73	-19.10	-19.24	-19.55	-18.39	-18.96
LFC 4394	0.2608	4.09 ± 0.33	15.37	-17.94	-19.84	-20.09	-20.33	-20.61	-19.56	-20.01
LFC 4536	0.2975	5.36 ± 0.88	16.88	-19.52	-20.46	-20.68	-21.02	-21.17	-20.18	-20.61
LFC 4763	0.0208	0.29 ± 0.01	1.84	-13.36	-13.86	-14.75	-15.22	-15.62	-13.27	-14.38

Note. — Column (1) gives our SExtractor catalog ID. Column (2) gives spectroscopic redshift obtained by us. Column (3) and (4) give Petrosian half light radius and semi-major axis length in kiloparsec. Columns (5) to (11) give absolute magnitudes of galaxies in SDSS filters. Column (10) and (11) give Johnson B and V absolute magnitudes obtained using Equations 3 and 4.

In their seminal papers on the isophotal shapes of elliptical galaxies, Bender et al. (1988) and Bender et al. (1989) define structural parameter a_n/a and b_n/a . The A_n and B_n provided by the `ellipse` table are normalized to the semi-major axis a and the local intensity gradient intensity $\frac{dI}{da}$ (`ellipse` output GRAD). The normalized B_n coefficients can be converted to the a_n/a parameter as

$$\frac{a_n}{a} = B_n \sqrt{1 - \epsilon} = B_n \sqrt{b/a}, \quad (6)$$

where b is the semi-minor axis and ϵ is isophotal ellipticity (Milvang-Jensen & Jørgensen 1999; Tremblay et al. 2007; Hao et al. 2006b). The factor of $\sqrt{1 - \epsilon} = \sqrt{b/a}$ is needed to re-normalize the radial deviation to the equivalent radius $r = \sqrt{ab}$ (Milvang-Jensen & Jørgensen 1999). The a_4/a parameter, which is the most dominant coefficient for any isophote deviating from a pure ellipse, quantifies the deviation along the major axis of the isophote. Isophotes with $a_4/a < 0$ have a “boxy” shape while the shape of isophotes with $a_4/a > 0$ is “disky” (Bender et al. 1988). The parameter b_n/a can be related to the normalized A_n parameter through a relation similar to Equation 6 but we do not use it in our present work.

The Fourier coefficients and morphological parameters vary along the semi-major axis, and we find that there are significant changes in values in the outer regions of the galaxies. It is therefore not enough to consider characteristic values of the parameters at a single fiducial distance from the centre, as has been done by earlier workers (see e.g. Bender et al. (1988), Hao et al. (2006)), for investigating relationships between these parameters and other global galaxy properties. However, in comparing data for a large sample of galaxies it is convenient to derive mean values of isophote parameters in a few regions, rather than the full sets of isophotes. We therefore divide each galaxy into four regions defined by distance from the centre along the semi-major axis: (i) seeing radius r_s to $1.5R_{50}$ (Region 1), (ii) $1.5R_{50}$ to $3.0R_{50}$ (Region 2), (iii) $3.0R_{50}$ to $4.5R_{50}$ (Region 3) and (iv) semi-major axis $> 4.5R_{50}$ (Region 4), where R_{50} is the Petrosian half-light radius. In each region we obtain the mean value of various parameters, weighted with the intensity (counts) and inversely weighted with the variance of the parameter. For example, the mean value of a_4/a in Region 1 is obtained as

$$\left\langle \frac{a_4}{a} \right\rangle = \frac{\int_{r_s}^{1.5R_{50}} \frac{a_4}{a}(r) I(r) \left[\sigma_{\frac{a_4}{a}}(r) \right]^{-2} dr}{\int_{r_s}^{1.5R_{50}} I(r) \left[\sigma_{\frac{a_4}{a}}(r) \right]^{-2} dr} \quad (7)$$

Note that not all galaxies have data in all four regions: for example, Region 1 does not exist if seeing radius $r_s > 1.5R_{50}$, while sometimes it is impossible to fit reliable ellipses in regions 3 and 4 due to crowding and/or low S/N . Some of the derived parameters for our sample of galaxies are listed in Table 3 (only the first 5 galaxies are listed here as examples; the complete table is at the end of the article as Table 10).

Table 3. Derived parameters for a sample of the LFC galaxies

ID (1)	B/T (2)	$\langle \frac{a_3}{a} \rangle$ (3)	$\langle \frac{a_4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)
LFC 462	0.89
		$3.52\text{e-}03 \pm 2.25\text{e-}03$	$-1.25\text{e-}03 \pm 2.16\text{e-}03$	0.057 ± 0.0058
		$7.01\text{e-}03 \pm 4.73\text{e-}03$	$6.91\text{e-}03 \pm 3.09\text{e-}03$	0.048 ± 0.0063
	
LFC 525	0.97
		$5.52\text{e-}03 \pm 1.89\text{e-}03$	$-1.09\text{e-}03 \pm 1.53\text{e-}03$	0.070 ± 0.0047
		$-6.61\text{e-}03 \pm 5.70\text{e-}03$	$1.21\text{e-}02 \pm 5.41\text{e-}03$	0.023 ± 0.0117
	
LFC 554	0.65	$-4.29\text{e-}03 \pm 1.96\text{e-}03$	$1.00\text{e-}02 \pm 1.82\text{e-}03$	0.094 ± 0.0040
		$-8.35\text{e-}03 \pm 3.77\text{e-}03$	$3.49\text{e-}03 \pm 3.75\text{e-}03$	0.115 ± 0.0071
	
	
LFC 558	0.68	$6.77\text{e-}03 \pm 1.47\text{e-}03$	$1.88\text{e-}03 \pm 1.42\text{e-}03$	0.038 ± 0.0035
		$4.92\text{e-}03 \pm 1.60\text{e-}03$	$1.12\text{e-}04 \pm 1.56\text{e-}03$	0.081 ± 0.0036
		$-1.10\text{e-}02 \pm 1.19\text{e-}02$	$-4.33\text{e-}03 \pm 1.28\text{e-}02$	0.164 ± 0.0248
	
LFC 622	0.59	$-4.88\text{e-}04 \pm 1.06\text{e-}03$	$1.51\text{e-}02 \pm 4.42\text{e-}04$	0.358 ± 0.0018
		$-1.18\text{e-}03 \pm 8.45\text{e-}04$	$1.29\text{e-}02 \pm 4.69\text{e-}04$	0.421 ± 0.0013
		$4.78\text{e-}02 \pm 4.35\text{e-}03$	$-1.47\text{e-}02 \pm 3.35\text{e-}03$	0.310 ± 0.0075
		$3.79\text{e-}02 \pm 1.57\text{e-}02$	$-2.56\text{e-}02 \pm 1.45\text{e-}02$	0.259 ± 0.0257
LFC 633	0.66	$5.30\text{e-}03 \pm 7.48\text{e-}04$	$-2.57\text{e-}03 \pm 6.74\text{e-}04$	0.213 ± 0.0016
		$1.59\text{e-}02 \pm 1.24\text{e-}03$	$-3.83\text{e-}03 \pm 1.11\text{e-}03$	0.219 ± 0.0027
		$1.64\text{e-}02 \pm 4.29\text{e-}03$	$-6.56\text{e-}04 \pm 4.22\text{e-}03$	0.269 ± 0.0074
		$-2.51\text{e-}01 \pm 7.91\text{e-}02$	$-2.39\text{e-}02 \pm 2.74\text{e-}02$	0.102 ± 0.0591

Note. — Here we list data for only the first five galaxies in our sample: the full table is at the end of the article as Table 10. Column (1) gives our SExtractor catalog ID. Column (2) gives bulge-to-total luminosity ratio (see Section 2.3). Columns (3) to (5) give average isophotal parameters a_3/a , a_4/a , and ellipticity in four different regions for each galaxy. Average values of isophotal parameters in Regions 1, 2, 3 and 4 are listed, when available, in the first, second, third and fourth rows respectively for each galaxy.

Our method of obtaining characteristic parameter values is similar to the method used by Hao et al. (2006) and Bender et al. (1988, 1989), except that (i) Hao et al. (2006) considered a single characteristic parameter value in the region $2r_s - 1.5R_{50}$ while Bender et al. (1988, 1989) defined the value in region $2r_s - 1.5r_e$, where r_e is the de Vaucouleur half light radius, and (ii) Bender et al. (1988, 1989) estimated the characteristic value of ellipticity as the maximum value in a peaked ellipticity profile, and as the value of the ellipticity at $1.5r_e$ in case of a continuously increasing or decreasing ellipticity profile. We have chosen multiples of R_{50} for defining the regions, as the value of r_e is very sensitive to the goodness of fit and the range of radius chosen for fitting surface brightness profiles with de Vaucouleurs law. The value of R_{50} has been obtained from the SDSS DR7 galaxy catalogue given by Blanton et al. (2005).

4. Observed properties of the galaxies

4.1. Some representative galaxies

In Figure 4 we show examples of three typical galaxies in our sample. The left panels show the observed isophotes (in blue) in each region of the galaxy along the semi-major axis, superimposed on grey-scale images from the LFC. The smooth red contours are the best-fitted ellipses to the observed contours (see Section 3). The right panels show the variation of a_4/a along the semi-major axis of the galaxies.

These three examples represent common patterns for galaxies in our sample. The a_4/a profile for the galaxy LFC1208_2549 shows diskiness (positive a_4/a) in all the regions except in Region 2. The diskiness of galaxy LFC1208_2549 peaks in Region 3. The profile of LFC1208_1585 shows boxiness (negative a_4/a) in outer regions and small diskiness in the inner regions. The profile for the galaxy LFC1208_4681 remains constant along the semi-major axis with value of a_4/a close to zero, as the isophotes of this galaxy do not deviate much from the elliptical shape. The vertical lines in black colour show the bins in which the mean values of isophotal parameters are computed. A red circle shows the mean value of a_4/a in each bin. The red dotted vertical line in each plot marks the seeing radius of the image frame. The region of each galaxy within the seeing radius is excluded from the study. The necessity of considering radial variation of isophotal parameters, as we do in this paper, is clear from the profiles in the figure.




Fig. 4.— The left panels show *i*-band image cut outs ($20'' \times 20''$) for three of our galaxies. Isophotes (in blue) are shown together with the best-fitted ellipses (in red). The right panels show the variation of a_4/a along the semi-major axis of each galaxy (with 1σ error bars). The dotted vertical lines in black colour indicate the bins of semi-major axis length in which the mean values of isophotal parameters are computed. A red circle shows the mean value of a_4/a in each bin. The positions of the blue contours in each left-hand panel are marked by blue crosses in the corresponding right-hand panels. The region of each galaxy within the seeing radius (dashed vertical line) is excluded from the study. Most previous studies of isophotal parameters have been confined to only the innermost of our four bins.

4.2. Sample properties

In this section we describe some of the basic properties of our sample and compare those with the properties of other samples used for similar studies. We look for evidence of systematic structural changes between the inner and outer regions of galaxies.

We first examine the differential probability distribution of ellipticity $\epsilon (= 1 - b/a)$, and shape parameters a_4/a and a_3/a in different regions of our sample galaxies.

We have fitted empirical Gaussian distribution functions (Equation 8) to the histograms of various shape parameters for our galaxies, in different radial regions. Although the intrinsic distributions are unknown *a priori*, empirical fits to the data may be useful for comparison between datasets or with theoretical studies. The results of the fits are given in Table 4 and the fits along with the distribution of the above parameters are shown in Figure 5. In some cases we could get better fits by fitting multiple Gaussians but the statistical significance and physical meaning of such fits was not clear. The differential probability distribution we use for ϵ , a_4/a and a_3/a are of the form

$$p(x)dx = [k_1 + k_2x + k_3e^{-\frac{1}{2}(\frac{x-\mu_1}{\sigma_1})^2}]dx$$

$$p(x)dx = [k_1 + k_2x + k_3e^{-\frac{1}{2}(\frac{x-\mu_1}{\sigma_1})^2} + k_4e^{-\frac{1}{2}(\frac{x-\mu_2}{\sigma_2})^2}]dx, \quad (8)$$

The coefficients k_3 , k_4 , μ_1 , μ_2 , σ_1 and σ_2 are the amplitude, mean and the standard deviation of the fitted Gaussian. The coefficients k_1 and k_2 are zero point and slope of the baseline for the Gaussian. The results of the fit are given in the Table 4 and the fits along with the distribution of the above parameters are shown in Figures 5 to 7.

The ellipticity distribution of the sample used by Hao et al. (2006) (dotted curve in Region 1, upper left panel of Figure 5) shows a peak around $\epsilon \sim 0.2$. The main peak in our sample, at $\epsilon \sim 0.1$, indicates the presence of a larger fraction of rounder galaxies. Both distributions of ellipticity drop to zero for $\epsilon > 0.5$. This difference in the ellipticity distributions is presumably because of the very different data sets. The galaxies of Hao et al. (2006) are all large and relatively nearby, with $z < 0.05$, selected over a large area of the sky. Most of our galaxies are much smaller and more distant, with a redshift distribution peaking at $z \sim 0.1$ and extending to beyond 0.5 (see Figure 2).

If Gaussian functions are fitted to the ellipticity distributions in Figure 5, we find that as we go from Region 2 to Region 4, the peak of the distribution occurs at around 0.13, 0.16 and 0.16, i.e. the peak shifts slightly towards more flattened ellipses. Overall, the distributions of ellipticity in Regions 1 and 2 are similar to each other, as are those for Regions 3 and 4, but there are significant differences between the inner and outer regions.

Table 4. Values of coefficients for fitted single and/or double Gaussians (see Equation 8 and Figures 5 to 7)

Parameter	k_1	k_2	k_3	μ_1	σ_1	k_4	μ_2	σ_2	χ^2	rms
ϵ (Region 1)	0.072 ± 0.056	-0.091 ± 0.131	0.158 ± 0.049	0.111 ± 0.012	0.057 ± 0.037	-	-	-	1.05	0.025
ϵ (Region 2)	0.036 ± 0.045	-0.041 ± 0.094	0.240 ± 0.040	0.127 ± 0.009	0.053 ± 0.013	-	-	-	1.45	0.023
ϵ (Region 3)	0.011 ± 0.014	0.0 ± 0.0	0.202 ± 0.017	0.159 ± 0.008	0.083 ± 0.012	-	-	-	1.03	0.022
ϵ (Region 4)	0.036 ± 0.018	0.0 ± 0.0	0.210 ± 0.032	0.160 ± 0.009	0.051 ± 0.020	-	-	-	1.01	0.033
$\frac{a_4}{a}$ (Region 1)	0.0 ± 0.0	0.0 ± 0.0	0.0486 ± 0.0152	0.0010 ± 0.0098	0.0131 ± 0.0053	0.3143 ± 0.0205	0.0022 ± 0.0001	0.0042 ± 0.0005	1.04	0.010
$\frac{a_4}{a}$ (Region 2)	0.0064 ± 0.3322	-0.0726 ± 2.1186	0.5163 ± 0.09308	0.0003 ± 0.0007	0.0016 ± 0.0024	0.1256 ± 0.5300	0.0015 ± 0.2297	0.0093 ± 0.2068	1.14	0.013
$\frac{a_4}{a}$ (Region 3)	0.0149 ± 0.0040	-0.0273 ± 0.0300	0.0683 ± 0.0632	-0.0270 ± 0.0038	0.0011 ± 0.0002	0.1240 ± 0.0107	0.00001 ± 0.00002	0.0079 ± 0.0013	0.92	0.020
$\frac{a_4}{a}$ (Region 4)	0.0268 ± 0.0076	-0.0267 ± 0.1084	0.1141 ± 0.0810	-0.0256 ± 0.0321	0.0014 ± 0.0005	0.1407 ± 2.6219	0.0110 ± 0.0004	0.0030 ± 0.0010	0.53	0.021
$\frac{a_3}{a}$ (Region 1)	0.0 ± 0.0	0.0 ± 0.0	0.4510 ± 0.0153	-0.0009 ± 0.0003	0.0078 ± 0.0004	0.0135 ± 0.0068	-0.0112 ± 0.0234	0.0712 ± 0.0208	0.92	0.008
$\frac{a_3}{a}$ (Region 2)	0.0 ± 0.0	0.0 ± 0.0	0.3639 ± 0.0307	-0.0011 ± 0.0010	0.0105 ± 0.0009	-	-	-	1.38	0.023
$\frac{a_3}{a}$ (Region 3)	0.0 ± 0.0	0.0 ± 0.0	0.2253 ± 0.0112	-0.0024 ± 0.0008	0.0142 ± 0.0009	0.0593 ± 0.0322	0.0513 ± 0.0043	0.0140 ± 0.0056	1.01	0.012
$\frac{a_3}{a}$ (Region 4)	0.0 ± 0.0	0.0 ± 0.0	0.1081 ± 0.0131	0.0046 ± 0.0061	0.0352 ± 0.0062	0.0090 ± 0.0231	0.0291 ± 0.0302	0.0036 ± 0.0009	0.93	0.023

Figure 6 shows the distribution of the quadrupole parameter a_4/a for our sample. The distribution in Region 1 shows a slight excess of disk isophotes (positive a_4/a values), although not as strong as that found at similar radii by Hao et al. (2006) and shown by the blue curve. However, the distribution we find in Region 2 (upper right panel) does resemble that of Hao et al. (2006). As in the case of the ellipticity plots in Figure 5, the histograms for a_4/a in Regions 1 and 2 are similar to each other but quite different from those for Regions 3 and 4.

The a_3/a parameter quantifies the deviation from pure ellipse that occur along the observed isophote every 120° . Pasquali et al. (2007) have suggested that such deviations may be attributed to the presence of dust or clumps present within the galaxy. But the distribution of the dust can be irregular in the galaxy and can also give rise to higher order Fourier coefficients. It may be better to interpret a_3/a as a structural parameter, possibly indicating dynamical effects from galaxy interactions or mergers. Figure 7 shows the distribution of this parameter for our sample.

The distribution of a_3/a in Region 1 is reasonably fitted by a sum of two Gaussian functions with mean -0.0009 and -0.0112 with dispersions 0.0078 and 0.0712 . The value of χ^2 per degree of freedom for the fit is $\chi^2_\nu = 0.92$. In comparison, the distribution of the Hao et al. (2006) sample is reasonably fitted by a single Gaussian with zero mean and dispersion of 0.0032 . In our sample the peaks of distribution in Region 1 and Region 2 occur at negative values of the parameter a_3/a while one peak of the double Gaussian in Region 3 and both peaks of the double Gaussian in Region 4 occur at positive values of parameter a_3/a (see Table 4). It is not known as to which property of galaxy determines the sign of a_3/a parameter. The findings of Jog & Maybhate (2006) suggest that non zero values of a_3/a parameter along with the presence of lopsidedness in the inner regions of galaxies can be interpreted as a signature of dynamically unrelaxed behaviour in the inner regions of the galaxies. These authors have Fourier-analysed the central few kpc of advanced mergers of galaxies using images from the Two-Micron All-Sky Survey (2MASS) and have obtained amplitudes and phases of Fourier components $m = 1, 2, 3$ and 4 . Their analysis indicates that in the case of mergers, the amplitudes $A1$ and $A2$ (for $m=1$ and 2 respectively) dominate over $m=3$ and 4 , and $A3$ is important only when $A1$ is large. $A1$ denotes the amplitude for the lopsidedness and is an indicator of mass asymmetry measured with respect to the constant center (see Section 3.2 of their paper for details). The study of lopsidedness in our sample galaxies should reveal whether the peaks of the a_3/a distribution around non-zero values in the inner regions of our galaxies can be interpreted as a signature of dynamically unrelaxed inner regions. Though we do not understand the reason for the occurrence of peaks at positive values of a_3/a in Region 3 and in Region 4, we can say that the processes which affect isophotal shape are different in the inner and outer regions.

We next look for correlations between the values for various shape parameters for individual galaxies, when adjacent radial regions are compared.

Figure 8 shows that there is a strong correlation between the ellipticity parameters in the innermost Regions 1 and 2 (upper panel). However, there is a significant change going from Region 2 to Region 3 (lower left); while most of the higher ellipticity galaxies remain correlated (those with $\epsilon > 0.3$ in Regions 1 and 2), about half of the nearly circular galaxies become more flattened. The outer most parts (Regions 3 and 4, lower right panel) show more scatter and fewer galaxies have data in Region 4, but an overall correlation remains.

We show similar diagrams for the shape parameter a_4/a in Figure 9. The dominant feature is a large increase in the scatter of points between Regions 2 and 3, shown in the lower right panel. The other two plots, for the outer Regions 3 and 4 (upper panel) and the inner Regions 1 and 2 (lower left) show some correlation between adjacent regions, with few points populating the top left and bottom right quadrants.

4.3. Are distribution of isophotal parameters statistically different in different regions?

We performed a two-sample Kolmogorov-Smirnov test (K-S test) to investigate whether the distribution of a_4/a , a_3/a and ϵ are same in our four regions. (see Tables 5, 6 and 7).

We used AstroStat⁴, which uses a public-domain statistical computing package *R*⁵ for statistical analysis, to perform the K-S test, in which the null hypothesis that the two samples belong to the same parent distribution is tested. The strength of evidence in support of a null hypothesis is given by probability P in the table. If the P is less than the assumed *level of significance* the hypothesis is rejected, where the *level of significance* is the probability of making a decision to reject the null hypothesis, when the null hypothesis is true.

We find from the K-S test that the distribution of ϵ in neighboring Regions belongs to the same parent distribution, as the test is accepted at the 5% level of significance. The distribution of a_4/a and a_3/a in Region 2 and Region 3 may not belong to the same parent distribution, as the test is rejected at the 5% level of significance. The distribution of a_4/a and a_3/a parameter in Region 1 and Region 2 are from same parent population, as suggested by high *P-value* ($=0.86$ and 0.61) obtained from the K-S test. Similarly, the distribution of

⁴<http://vo.iucaa.ernet.in/~voi/AstroStat.html>

⁵<http://www.r-project.org/>

Table 5. Results of two sample K-S test for the a_4/a parameter

Sample	Region 1		Region 2		Region 3		Region 4	
	D^1	P^2	D	P	D	P	D	P
Hao et al. (2006)	0.15	0.041	0.13	0.038	0.22	5.97e-04	0.29	5.81e-04
Region 1			0.08	0.855	0.25	0.005	0.35	6.41e-04
Region 2					0.21	0.014	0.33	5.44e-04
Region 3							0.16	0.383

¹The K-S statistic D is defined as the maximum value of the absolute difference between two cumulative distribution functions where the cumulative distribution function is obtained from the list of data points of each sample on which K-S test is applied.

² P gives the level of significance with which the null hypothesis may be accepted. Small values of P imply that the cumulative distribution function of two samples tested are significantly different (see Press et al. 1992). The confidence that both the population do not belong to the same parent distribution is given by $(1 - P) \times 100$.

Table 6. Results of two sample K-S test for the a_3/a parameter ¹

Sample	Region 1		Region 2		Region 3		Region 4	
	D	P	D	P	D	P	D	P
Hao et al. (2006)	0.30	2.85e-07	0.30	1.65e-09	0.33	1.16e-08	0.44	1.34e-08
Region 1			0.10	0.613	0.21	0.031	0.29	0.007
Region 2					0.19	0.034	0.28	0.005
Region 3							0.19	0.151

¹Notation as in table Table 5.

a_4/a and a_3/a parameter in Region 3 and Region 4 are same as the test is accepted at the 5% level of significance, though the strength of evidence in support of the null hypothesis is not as strong as it is for Region 1 and Region 2. The Results of K-S test clearly shows the discontinuity in the distribution of parameters as we go from Region 2 to Region 3.

4.4. Frequency of boxy and disk ellipticals

Bender et al. (1989) found that $\sim 1/3$ of their sample galaxies show boxy isophotes, $\sim 1/3$ pointed isophotes and $\sim 1/3$ of isophotes have deviation smaller than 0.2% of the semi-major axis length. The frequency of boxy and disk isophotes in different radial regions of our sample galaxies is given in Table 8. We find a larger fraction of disk isophotes in Regions 1 and 2. The diskiness is generally attributed to the presence of a weak edge-on disk superposed on the spheroidal main body. One of the possible reasons for the increased fraction of disk isophotes in Region 1 is detection of those weak disks in our sample galaxies which would not be detected with relatively low SNR images.

The major fraction of our galaxies have either boxy or disk isophotes in their outer regions: we find that the fraction of near-circular galaxies having $|a_4/a| \leq 0.2\%$ is less than 10% in Regions 3 and 4, compared with 25% to 30% in Regions 1 and 2. This is consistent with the expectation that tidal extensions and other environmental effects are likely to be stronger in the outer regions as compared to the inner regions of galaxies.

We also find a higher frequency of boxy as compared to disk isophotes in the outer regions. It has been suggested by Nieto & Bender (1989) that tidal extensions may also cause pointed isophotes. The higher fraction of boxy isophotes indicates that: (i) the presence of additional forces along with the tidal extensions; and (ii) that such forces are either more frequent or more effective in outer regions.

The frequency of disk isophotes is higher in the outer regions of dwarf early-type galaxies having $M_B > 17.0$ in our sample. The fractions of disk isophotes for such galaxies in Region 1, Region 2, Region 3 and Region 4 are 1/6, 4/8, 4/6 and 2/2 respectively. The denominators of above fractions for each region are total number of dwarf early-type galaxies for which isophotal parameters are available. This higher fraction of disk isophotes in outer regions from Region 1 to Region 4 observed for dwarf early-type galaxies in our sample is quite different from the corresponding frequency in different regions when we consider all the galaxies in the sample. Differences in the environment of the dwarf and luminous early-type galaxies may explain the observed differences of their isophotal properties, as dwarf elliptical galaxies are found in high-density regions, either in galaxy clusters or in locations near more

Table 7. Results of two sample K-S test for the $(1 - b/a)$ parameter ¹

Sample	Region 1		Region 2		Region 3		Region 4	
	D	P	D	P	D	P	D	P
Hao et al. (2006)	0.28	3.31e-06	0.28	3.93e-08	0.20	0.002	0.14	0.30
Region 1			0.11	0.466	0.19	0.068	0.27	0.002
Region 2					0.16	0.098	0.23	0.039
Region 3							0.13	0.574

¹Notation as in table Table 5.

Table 8. Frequency of boxy and diskly isophotes

Radial bin	Region	Number of galaxies with data in region	boxy	diskly	$ a_4/a \leq 0.2\%$
$r_s - 1.5R_{50}$	Region 1	96	25 %	48%	27%
$1.5R_{50} - 3.0R_{50}$	Region 2	131	28%	43%	29%
$3.0R_{50} - 4.5R_{50}$	Region 3	96	55%	36%	09%
$r > 4.5R_{50}$	Region 4	52	54%	40%	06%

massive spiral and elliptical galaxies (Fergusone & Sandage 1989; Van et al. 2004).

4.5. Correlation between isophotal shape and ellipticity

Figure 10 shows plots of a_4/a against ellipticity ϵ for galaxies in each of our four radial regions. Our data are most complete for Region 2; we miss some galaxies in Region 1 when the ‘seeing’ $r_s > 1.5R_{50}$, and from Regions 3 and 4 due to confusion and low S/N. The solid lines in each panel define a chevron-shaped region within which all galaxies from the sample of Bender et al. (1989) were found. The dotted lines are extrapolations of these lines to higher values of ellipticity.

The distributions in Regions 1 and 2 are similar to each other, and most galaxies lie within the zone defined by Bender et al. (1989). Moreover, our galaxies show the same trend, with a preponderance of disk-like isophotes (with positive a_4/a) which extend to larger values as the ellipticity increases. However, we do find some galaxies lying outside Bender et al. (1989)’s boundaries, in particular a group with boxy isophotes (negative a_4/a) and low ellipticity in Region 2. We also find some galaxies with large ellipticity ($1 - b/a > 0.4$) but with a_4/a near zero.

Our data become increasingly incomplete and with larger errors in Regions 3 and 4, where we have plotted only points with errors less than 0.01 and 0.015 respectively. The pattern appears similar in the two lower panels of Figure 10, and different from the pattern in the inner regions, although this may be at least partly due to the increasing errors and smaller samples.

Overall, the isophote shape data are consistent with the patterns seen in Section 4.4 and clearly indicate that the phenomena responsible for keeping the isophotes close to an elliptical shape at small radii (of the rounder galaxies) weaken as we go further out along the radius. Another possibility is that the effects of tidal interaction become more dominant, to make the isophotes deviate from the elliptical shape. Environmental studies may show whether this is true.

5. Discussion

5.1. Are projection effects important?

The results obtained by Bender et al. (1989) indicate beyond doubt that the boxiness or diskiness is an intrinsic property of the early-type galaxies rather than an effect caused

by projection. This means that the sign of the a_4/a parameter is independent of projection effects, but it is not possible to say how the absolute value of a_4/a depends on the viewing angle. The trend found by Bender et al. (1989) and Hao et al. (2006), that the larger values of $|a_4/a|$ are observed for galaxies which appear more elliptical is consistent, with a few exceptions, with the results obtained for our sample galaxies in Section 4.5.

5.2. Comparison with N-body simulations

The differences in the properties of boxy and disk elliptical are also supported by the N-body simulations of major mergers (e.g. Hernquist 1993; Lima Neto & Combes 1995; Khochfar & Burkert 2005), but some of the findings of N-body simulations are not consistent with the observations. For example, Stiavelli et al. (1991) and Heyl et al. (1994) analyzed the isophotal shapes of the remnants from their N-body simulations of dissipationless collapse, and found that the same object has value of a_4 negative or positive depending on the viewing angle. Similarly, Fabio et al. (1993) have reported that isophotal shape can not discriminate between the possible origin of the early-type galaxies, at least in the range $-1 \leq a_4/a \times 100 \leq 1$ because of the dependence of the shape upon the viewing angle as seen using simulations. We do not understand the dependence of a_4/a parameter on viewing angle found by above authors which is inconsistent with the observations. In future, the understanding of true intrinsic shapes of early-type galaxies may elucidate the effect of projection on the value of a_4/a parameter.

N-body merger simulations have been widely used by various researchers to study the origin of boxy and disk ellipticals. Simulations have been done taking progenitors for the merger as (i) a combination of galaxies with different morphologies (ii) a combination of galaxies with different mass ratios. Bournaud et al. (2005) have shown in their Fig. 2, that a 7:1 merger produces a galaxy with boxy-isophotes in the inner region while the outer region is highly disk. Such radial variation has also been reported using the analysis of 2MASS data for the Arp mergers (see Chitre & Jog 2002, Appendix). Similar merger scenarios can be considered as possible origin of the inner boxy and outer disk isophotes observed in some of our sample galaxies.

Bournaud et al. (2005) have shown that for a merger with 7:1 mass ratio, outer diskiness is observed for a $25 \text{ mag arcsec}^{-2}$ isophote around 20 kpc from the centre (see their Fig. 2 & 3). In the region $\sim 20 - 30$ kpc, the surface density falls by a factor of $\sim 80 - 100$ and for a constant mass-to-light ratio this corresponds to a $\sim 4 - 5$ magnitude difference. The outer region of Bournaud et al. (2005) corresponds to the Region 4 of most of our sample galaxies, thus a disk is clearly indicated at such large radii.

A typical value of $a_4/a \simeq 0.01 - 0.02$ is obtained for the remnants of collisionless N-body simulations of binary mergers of disk galaxies with mass ratios of 1:1, 2:1, 3:1 and 4:1 (see Nabb & Burkert 2003, Fig. 3), while observations show larger diskiness in early-type galaxies. For example, observed values $a_4/a \lesssim 0.03$ are obtained for the galaxies studied by Bender et al. (1989), while some of our sample galaxies show larger diskiness ($a_4/a > 0.03$). Such galaxies are also a part of sample studied by the Hao et al. (2006b, Fig. 1). The origin of strong diskiness therefore should have a different merger scenario from that studied by Nabb & Burkert (2003). The galaxies with larger diskiness could well be explained by hybrid mergers, in mass ratio 4:1-10:1 (Bournaud et al. 2004, 2005). They report $a_4/a = 0.064 \pm 0.01$ for mergers with mass ratio 4.5:1 and $a_4/a \simeq 0.07$ for mergers with mass ratio 7:1.

In binary merger scenarios, disky ellipticals are produced by 3:1-4:1 mergers while boxy ellipticals are product of equal-mass 1:1 mergers (Bournaud et al. 2007). But equal-mass binary mergers fail to reproduce the most boxy elliptical galaxies, particularly giant ellipticals (Nabb & Burkert 2003; Nabb & Ostriker 2009). Alternatively, multiple minor mergers can reproduce boxy ellipticals, where the product of mergers mainly depends on the total merged mass. When total merged mass increases (≥ 1.6), on average the remnants tend to show boxy isophotes (see Bournaud et al. 2007, Fig. 5). The merged mass has been defined in a particular way by authors in their paper: for instance, when the initial galaxy has merged with 3 companions, each of them having a 5:1 mass ratio, this so-called merged mass is 1.6 (1.0 for the main initial galaxy and 0.2 for each companion). Figure 11 shows a plot of a_4/a with absolute i -band magnitude for the galaxies with redshift information. We find that a larger number of galaxies with boxiness have $M_i < -19.5$, in all regions along the radius. Since more luminous galaxies will be more massive, the above trend can be considered similar to the trend observed for boxiness of the remnants of multiple minor mergers observed by Bournaud et al. (2007), though we do not exactly know the mass corresponding to $M_i < -19.5$.

We have tried to compare our observations with the isophotal shapes of the merger remnants, but it is not in the scope of this paper to trace the merger history of the sample galaxies on the basis of observed isophotal shapes.

6. Conclusions

In this paper, we have studied the properties of the isophotal shapes of early-type galaxies (E/SO) to very faint outer regions, well beyond the levels reached by previous studies of this type. Our study is based on deep 5-color Hale Telescope LFC CCD images of several fields, originally taken for a different project, that enable us to reach surface

brightnesses some $4 \text{ mag arcsec}^{-2}$ fainter than most similar studies. Here we present the first results for one $14' \times 25'$ field.

We selected target galaxies using the deep LFC *i*-band image of the field and used much shallower SDSS imaging data to get basic parameters such as apparent magnitudes and Petrosian radii for each galaxy. 266 sufficiently large and bright galaxies were selected for further study, 132 of which were identified as being of early-type on the basis of their bulge-to-total light ratios. We obtained spectra for over half of this sample using the multi-fibre system AAOmega on the AAT. These yielded reliable redshifts for 53 of the early-type galaxies, enabling us to derive their absolute magnitudes and physical sizes.

We fitted a sequence of ellipses to successive isophotes in the deep LFC images and derived a range of isophotal shape parameters that measure their ellipticity and orientation, and also higher order departures from a purely elliptical shape. We then derive mean values for these parameters in four radial bins along the semi major axis of each galaxy. We find empirical fitting formulae for the probability distribution of the different isophotal parameters in each bin, which will be useful for comparison with theoretical studies, e.g. from N-body simulations.

Finally, we have investigated possible correlations of isophotal shape parameters with other global properties of the galaxies, and inspected whether the correlations change along the radius. We find that the isophotal shapes of the inner regions of our sample of galaxies are statistically different from the isophotal shapes observed in the outer regions. In the central regions we see patterns similar to those seen in previous studies of nearby galaxies, with some galaxies showing 'boxy' isophotes while others appear 'disky'. However, the pattern seen in the inner region of each galaxy tends to change as the radius increases, suggesting that while the inner parts of the galaxies are coherent and presumably the result of specific dynamical processes, at larger radii the shapes and orientations of the isophotes change and the behavior is not well-defined. This may indicate effects from the formation and evolution of each galaxy which are not yet fully relaxed. However, for a full analysis of these data we need to know the distances to the galaxies so that we can derive their luminosities, actual sizes and other physical parameters. We hope to obtain fibre spectra for many more galaxies to do this, preferably with one of the new Integral Field Unit (IFU) systems that would avoid the effects of 'aperture bias' that arise in single fibre spectra, and enable us to compare galaxies at very different redshifts."

7. Acknowledgments

Two of the authors (LC and SKP) are grateful to ISRO for providing funds for this project under their RESPOND scheme (project no. ISRO/RES/2/343/2007-08). A.M. was supported by NSF grants AST-0407488 and AST-0909182. This paper makes use of deep LFC data obtained at the California Institute of Technologys Palomar Observatories and we acknowledge the role of S.G. Djorgovski, M. Bogosavljevic, and E. Glikman that resulted in the data Facility: Hale Telescope. We are thankful to Prof. Chanda Jog for giving valuable suggestions through email communication regarding comparison of our work with results from N-body simulations.

We thank the anonymous referee for valuable comments.

We thank the staff of the Australian Astronomical Observatory for obtaining the AAT AAOmega spectra for us during Service Observing.

Funding for the Sloan Digital Sky Survey (SDSS) and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, and the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium (ARC) for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, The University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

A. Estimation of surface brightness limit with LFC image as compared to the SDSS image

Following is the estimation of how far in the surface brightness we can go with LFC images as compared to SDSS images assuming the poisson statistics.

The number of source and background photons collected in t seconds using a telescope of diameter D is

$$\begin{aligned} N_s &= n_s D^2 t \\ N_b &= n_b D^2 t \end{aligned} \quad (\text{A1})$$

where n_s and n_b are rates of photons received in unit area from source and the background. Signal-to-noise ratio (SNR) for an extended source which spans a solid angle Ω can be given as

$$\frac{N_s}{\sigma_s} = \frac{n_s (\Omega D^2 t)^{\frac{1}{2}}}{\sqrt{n_s + 2n_b}} \quad (\text{A2})$$

For $n_s \ll n_b$, the above equation becomes

$$\frac{N_s}{\sigma_s} = \frac{n_s (\Omega D^2 t)^{\frac{1}{2}}}{\sqrt{2n_b}} \quad (\text{A3})$$

The SNR will increase by factor of \sqrt{P} , if the source intensity is obtained by averaging intensity in P number of pixels,

$$SNR_{av} = \frac{N_s \sqrt{P}}{\sigma_s} \quad (\text{A4})$$

In ellipse fitting task to get the average intensity of an isophotes the intensity is averaged in elliptical annulus along the fitted ellipse. The output table of **ellipse** gives parameters INTENS and INT_ERR. Where the INTENS is the mean intensity along the fitted ellipse and the INT_ERR is the total error in the intensity. We can write

$$\frac{INTENS}{INT_ERR} = \frac{N_s \sqrt{P}}{\sigma_s} \quad (\text{A5})$$

If the isophotes in two different images have same $\frac{INTENS}{INT_ERR}$ then

$$\frac{n_{s1} (P_1 \Omega_1 D_1^2 t_1)^{\frac{1}{2}}}{\sqrt{2n_{b1}}} = \frac{n_{s2} (P_2 \Omega_2 D_2^2 t_2)^{\frac{1}{2}}}{\sqrt{2n_{b2}}} \quad (\text{A6})$$

The difference in surface brightness of isophotes ($\mu_{s1} - \mu_{s2}$) with same $\frac{INTENS}{INT_ERR}$ in two different images can be given as

$$\mu_{s1} - \mu_{s2} = \frac{1}{2} \left[\mu_{b1} - \mu_{b2} - 2.5 \log \left[\left(\frac{D_2}{D_1} \right)^2 \left(\frac{t_2}{t_1} \right) \left(\frac{\Omega_2}{\Omega_1} \right) \left(\frac{P_2}{P_1} \right) \right] \right] \quad (\text{A7})$$

The ellipse is fitted till $\frac{INTENS}{INT_ERR}$ drops to 3 to derive surface brightness profiles in the images of SDSS and LFC (see Figure 1 as an example). Putting Values for SDSS image: $D_1=2.5\text{m}$, $t_1=54\text{s}$, $P_1=72$ pixels, $\mu_{b1} = 20.3 \text{ mag arcsec}^{-2}$ and for LFC image: $D_1=5\text{m}$, $t_1=9065\text{s}$, $P_1=184$ pixels, $\mu_{b1} = 20.2 \text{ mag arcsec}^{-2}$, in Equation A7 we expect to go minimum of $\sim 4 \text{ mag arcsec}^{-2}$ deeper in the LFC image as compared to the SDSS image.

REFERENCES

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Bailey J. A. et al., 2004, The 2DFDR Data Reduction System Users Manual. AAO (<http://www.aao.gov.au/AAO/2dF/manual.html>)
- Bender, R., Döbereiner, S., Möllenhoff, C. 1988, *A&AS*, 74, 385
- Bender, R., Surma, P., Döbereiner, S., Möllenhoff, C., Madejsky, R. 1989, *A&A*, 217, 35
- Bertin, E. & Arnouts, S. 1996, *A&AS*, 117, 393
- Blanton, M.R. et al. 2005, *AJ*, 129, 2562
- Blanton, M.R. & Roweis, S. 2007, *AJ*, 133, 734
- van den Bosch, F. C. 1994, *AJ*, 108, 1579
- Bournaud, F., Combes F., Jog, C. J. 2004, *A&A*, 418, 27
- Bournaud, F., Jog, C. J., Combes F. 2005, *A&A*, 437, 69
- Bournaud, F., Jog, C. J., Combes F. 2007, *A&A*, 476, 1179
- Busko, I. C. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 139
- Cannon, R. D., et al. 2006, *MNRAS*, 372, 425
- Capaccioli, M., Piotto, G., Rampazzo, R., 1988, *AJ*, 96, 487
- Chitre, A & Jog, C. J. 2002, *A&A*, 388, 407
- Fabio G., Luca, R., Roberto, R. 1993, *MNRAS*, 261, 379
- Ferguson, H. C., & Sandage, A. 1989, *ApJ*, 346, L53
- Ferrarese, L., van den Bosch, F.C., Ford, H.C., Jaffe, W., O’Connell, R.W. 1994, *AJ*, 108, 1598
- Freeman, K. C. 1970, *ApJ*, 160, 811
- Hao, C.N., Mao, S., Deng, Z.G., Xia, X.Y., Wu, H. 2006, *MNRAS*, 370, 1339

- Hao, C.N., Mao, S., Deng, Z.G., Xia, X.Y., Wu, H. 2006, MNRAS, 373, 1264
- Hernquist, L. 1993, ApJ, 409, 548
- Heyl, J.S., Hernquist, L., Spergel, D.N. 1994 ApJ, 427, 165
- Jedrzejewski, R. 1987, MNRAS, 226, 747
- Jog, C. J., & Maybhate, A. 2006, MNRAS, 370, 891
- Khochfar, S., & Burkert A. 2005, MNRAS, 341, 33
- Lauer, T. R. 2005, AJ, 129, 2138
- Lima Neto, G. B., & Combes, F. 1995, A&A, 294, 657
- Mahabal, A. et al. 2005, ApJ, 634, L9
- Malin, D. & Carter, D., 1980, Nature, 285, 643
- Milvang-Jensen B., Jørgensen I. 1999, Baltic Astronomy, 8, 535
- Miszalski et al. 2006, MNRAS, 371, 1537
- Naab, T., Burkert, A., Hernquist, L. 1999, ApJ, 523, L133
- Naab, T., & Burkert, A. 2003, ApJ, 597, 893
- Naab, T., Khochfar, S., & Burkert, A. 2006, ApJ, 636, L81
- Naab, T., & Ostriker, J. P. 2009, ApJ, 690, 1452
- Nieto J. L. & Bender R. 1989, A&A, 215, 266
- Pasquali A., van den, Bosch, F. C., Rix, H. -W. 2007, ApJ, 664, 738
- Peng, C. Y., et al. 2002, AJ, 124, 266
- Poggianti, B. M. 1997, A&AS, 122, 399
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., & Flannery, B.P. 1992, Numerical Recipes in Fortran (Cambridge: Cambridge University Press), p. 614
- Quinn, P. 1984, ApJ, 279, 596
- Rawat, A., et al. 2007, A&A, 496, 483

- Rest, A. 2001, *AJ*, 121, 2431
- Ryden, B. S., Forbes, D. A., Terlevich, A. I. 2001, *MNRAS*, 326, 1141
- Saunders, W., Bridges, T., Gillingham, P., et al. 2004, *Proc. SPIE*, 5492, 389
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M., 1998, *ApJ*, 500, 525
- Sersic, J. L. 1968, *Atlas de Galaxias Australes* (Cordoba: Obs. Astron, Univ. Nac. Cordoba)
- Smith, J. A. et al. 2002, *AJ*123, 2121
- Stiavelli, M., Londrillo, P., Messina, A. 1991, *MNRAS*, 251, 57
- Tremblay, G. R. et al. 2007, *ApJ*, 666, 109
- Van Zee L., Skillman E. D., Haynes M. P. 2004, *AJ*, 128, 121
- Vincent, R. A. & Ryden B. S. 2005, *ApJ*, 623, 137




Fig. 5.— Distribution of ellipticity $\epsilon = (1 - b/a)$, in different regions. The dashed lines are the best fitted Gaussian distributions with baseline (see text and Table 4). The dotted curve in Region 1 is the best fit Gaussian to the ellipticity distribution obtained by Hao et al. (2006) for their sample. The error bars indicate 1σ Poisson errors.




Fig. 6.— Distribution of a_4/a parameters, in different regions. The dashed lines are best fitted single or double Gaussians, (see text and Table 3). The dotted curve in Region 1 is the best fit Gaussian to the a_4/a distribution obtained by Hao et al. (2006) for their sample. The error bars indicate 1σ Poisson errors.




Fig. 7.— Distribution of a_3/a parameters, in different regions. The dashed lines are best fitted single or double Gaussians (see text and Table 4). The dotted curve in Region 1 is the best fit Gaussian to the a_3/a distribution obtained by Hao et al. (2006) for their sample. The error bars indicate 1σ Poisson errors.




Fig. 8.— Correlation of ellipticity ($\epsilon = 1 - b/a$) parameters from neighboring regions. The median error bars are shown at the the top right-hand corner in each pannel.




Fig. 9.— The correlation of a_4/a parameters from neighboring regions. The error bars are as in Figure 8.




Fig. 10.— Isophote shape parameter a_4/a plotted against ellipticity, in the four radial regions. In Regions 3 and 4 we omit galaxies with errors $\sigma(a_4/a) > 0.01$ and 0.015 respectively, for clarity. Bender et al. (1989) found that their galaxies lay within the chevron-shaped region between the pairs of solid diagonal lines.




Fig. 11.— Isophotal shape parameter a_4/a plotted against absolute Petrosian i magnitude.

Table 9. Basic parameters of the sample galaxies

Our ID	SDSS ID	RA	Dec	u mag	g mag	r mag	i mag	z mag	R_{50} arcsec	sma arcsec	μ_i mag arcsec ⁻²	SNR
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
LFC 462	588848899913548358	182.02545140	-0.00014042	21.50 ± 0.43	21.06 ± 0.13	20.55 ± 0.13	20.57 ± 0.21	19.96 ± 0.43	1.07 ± 0.16	3.81	26.36 ± 0.06	17.25
LFC 525	587748929241612656	182.10068456	0.00579127	22.48 ± 1.58	20.76 ± 0.09	20.42 ± 0.10	20.13 ± 0.12	20.74 ± 0.95	1.10 ± 0.11	3.81	26.06 ± 0.04	26.07
LFC 554	587748929241547219	181.99182097	0.00889758	22.09 ± 1.04	21.89 ± 0.24	21.11 ± 0.18	19.90 ± 0.09	19.67 ± 0.34	1.28 ± 0.07	3.81	26.67 ± 0.08	12.50
LFC 558	587748929241547253	181.99959155	0.01032991	26.27 ± 2.14	21.73 ± 0.24	20.20 ± 0.09	19.44 ± 0.07	19.05 ± 0.21	1.01 ± 0.07	5.11	26.58 ± 0.10	10.06
LFC 622	587748929241546923	182.03581581	0.01761600	20.79 ± 0.41	19.26 ± 0.03	18.16 ± 0.02	17.70 ± 0.02	17.33 ± 0.06	1.38 ± 0.02	10.11	27.21 ± 0.09	11.88
LFC 633	587748929241547046	182.04195384	0.01802068	20.48 ± 0.30	20.09 ± 0.06	19.51 ± 0.05	19.11 ± 0.05	19.21 ± 0.28	1.26 ± 0.05	8.32	27.18 ± 0.08	13.77
LFC 663	587748929241612845	182.171111251	0.01778466	21.46 ± 0.60	21.97 ± 0.26	20.89 ± 0.15	20.06 ± 0.10	19.19 ± 0.22	1.14 ± 0.08	3.81	25.90 ± 0.05	23.48
LFC 677	587748929241547621	181.99102394	0.02035316	19.94 ± 0.20	22.14 ± 0.42	20.94 ± 0.16	20.98 ± 0.33	22.79 ± 4.34	0.80 ± 0.17	5.36	27.64 ± 0.19	5.11
LFC 725	587722983351583067	182.01578218	0.02274361	25.14 ± 4.14	21.58 ± 0.27	20.82 ± 0.13	20.75 ± 0.30	20.12 ± 0.77	1.32 ± 0.37	3.81	26.65 ± 0.12	8.92
LFC 741	587748929241547216	181.99078515	0.02434438	21.46 ± 0.57	21.65 ± 0.19	20.74 ± 0.12	20.19 ± 0.12	19.80 ± 0.36	0.97 ± 0.09	4.86	27.60 ± 0.21	4.73
LFC 824	587748929241547430	182.05600812	0.03170720	23.94 ± 10.03	23.14 ± 2.52	20.28 ± 0.18	20.44 ± 0.65	19.60 ± 0.78	0.74 ± 0.33	3.81	25.93 ± 0.04	28.48
LFC 848	587748929241547282	182.00945210	0.03522617	22.05 ± 1.24	20.93 ± 0.14	20.26 ± 0.11	20.13 ± 0.17	19.63 ± 0.39	1.00 ± 0.12	3.81	26.89 ± 0.09	11.38
LFC 925	587748929241612771	182.14531708	0.03984272	21.09 ± 0.52	21.00 ± 0.13	20.09 ± 0.07	19.76 ± 0.09	19.08 ± 0.24	1.27 ± 0.07	3.81	26.10 ± 0.05	21.72
LFC 954	587748929241612750	182.13801353	0.04149656	24.88 ± 5.75	23.04 ± 0.87	20.93 ± 0.17	20.11 ± 0.14	20.28 ± 0.78	1.06 ± 0.15	3.63	26.12 ± 0.08	12.67
LFC 985	587748929241547693	182.03813760	0.04718642	25.96 ± 2.24	25.08 ± 1.99	20.94 ± 0.14	19.70 ± 0.08	19.39 ± 0.25	1.21 ± 0.10	6.21	28.07 ± 0.20	4.93
LFC 988	587748929241547668	182.02239309	0.04442920	23.62 ± 2.84	22.52 ± 0.36	21.55 ± 0.20	20.59 ± 0.13	20.09 ± 0.37	0.79 ± 0.09	3.81	26.41 ± 0.06	18.00
LFC 1013	587748929241547396	182.04304159	0.04740958	21.50 ± 0.78	21.27 ± 0.18	20.58 ± 0.11	20.18 ± 0.15	19.32 ± 0.32	1.36 ± 0.22	3.81	25.31 ± 0.02	61.15
LFC 1039	587748929241612867	182.17918290	0.04867795	22.04 ± 1.41	21.40 ± 0.21	20.46 ± 0.10	20.72 ± 0.26	22.75 ± 4.14	0.93 ± 0.17	5.91	27.94 ± 0.24	4.04
LFC 1042	587748929241547353	182.02923093	0.04940430	23.60 ± 2.57	22.65 ± 0.35	21.28 ± 0.15	20.49 ± 0.11	20.85 ± 0.70	0.77 ± 0.07	4.86	26.97 ± 0.11	9.38
LFC 1044	587748929241546926	182.04119522	0.05269923	21.45 ± 0.79	19.67 ± 0.04	19.09 ± 0.04	18.87 ± 0.05	18.40 ± 0.15	1.40 ± 0.06	8.74	28.13 ± 0.24	3.96
LFC 1056	587748929241612449	182.15112952	0.05083776	21.45 ± 0.69	19.87 ± 0.04	19.31 ± 0.04	19.01 ± 0.05	19.04 ± 0.23	1.20 ± 0.05	7.92	27.60 ± 0.12	8.37
LFC 1091	587748929241547628	181.99519742	0.05510190	24.95 ± 7.55	22.85 ± 0.91	20.78 ± 0.13	20.12 ± 0.17	19.46 ± 0.45	1.02 ± 0.16	5.11	27.28 ± 0.15	6.76
LFC 1124	587748929241547307	182.01715327	0.05830780	25.45 ± 2.61	22.96 ± 0.45	21.02 ± 0.12	19.92 ± 0.06	19.36 ± 0.18	0.91 ± 0.04	3.81	26.17 ± 0.04	23.85
LFC 1204	587748929241612881	182.18147441	0.07250930	21.71 ± 0.83	22.33 ± 0.41	20.33 ± 0.11	19.78 ± 0.10	19.52 ± 0.33	1.37 ± 0.14	6.21	28.01 ± 0.30	3.08
LFC 1239	587748929241612532	182.10134588	0.06863121	23.23 ± 2.07	20.65 ± 0.06	19.37 ± 0.03	18.94 ± 0.03	18.88 ± 0.13	0.74 ± 0.02	6.52	26.94 ± 0.08	12.72
LFC 1263	587748929241547309	182.01728221	0.06837554	21.55 ± 0.42	22.33 ± 0.24	21.41 ± 0.16	20.89 ± 0.14	21.69 ± 1.29	0.78 ± 0.07	3.63	26.63 ± 0.10	10.34
LFC 1273	587748929241612910	182.18810764	0.06914050	21.82 ± 0.53	21.66 ± 0.13	20.84 ± 0.09	20.17 ± 0.07	20.00 ± 0.30	0.87 ± 0.05	3.81	27.01 ± 0.12	8.71
LFC 1308	587748929241612921	182.19052619	0.07268490	22.26 ± 1.23	21.19 ± 0.13	20.13 ± 0.07	19.72 ± 0.08	19.09 ± 0.20	1.06 ± 0.06	4.63	26.72 ± 0.10	10.47
LFC 1369	587748929241612903	182.18741821	0.07738771	21.31 ± 0.52	21.93 ± 0.25	21.04 ± 0.17	20.26 ± 0.12	19.98 ± 0.46	1.06 ± 0.09	3.81	26.61 ± 0.07	14.24
LFC 1385	587748929241612551	182.19799673	0.08086235	22.12 ± 0.99	20.31 ± 0.07	19.38 ± 0.05	18.93 ± 0.05	18.77 ± 0.14	0.97 ± 0.05	3.81	24.98 ± 0.02	58.15
LFC 1411	587748929241612876	182.18108286	0.08007677	21.93 ± 0.92	21.36 ± 0.15	20.44 ± 0.10	20.25 ± 0.12	20.21 ± 0.56	1.02 ± 0.09	3.81	27.26 ± 0.17	6.08
LFC 1437	587748929241547418	182.04924891	0.08279051	25.63 ± 3.39	21.95 ± 0.25	21.11 ± 0.17	20.90 ± 0.24	20.20 ± 0.53	0.86 ± 0.15	3.81	26.67 ± 0.08	13.20
LFC 1456	587748929241547286	182.01007203	0.08412366	23.59 ± 3.40	22.08 ± 0.27	20.79 ± 0.14	20.04 ± 0.10	19.17 ± 0.21	0.93 ± 0.08	3.81	26.48 ± 0.06	16.67

Table 9—Continued

Our ID	SDSS ID	RA	Dec	u mag	g mag	r mag	i mag	z mag	R_{50} arcsec	sma arcsec	μ_i mag arcsec ⁻²	SNR
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
LFC 1459	587748929241612905	182.18630107	0.08511143	21.18 ± 0.51	21.35 ± 0.16	20.11 ± 0.07	19.73 ± 0.08	19.40 ± 0.29	1.21 ± 0.06	5.63	27.34 ± 0.18	5.42
LFC 1477	587748929241547320	182.01899895	0.08676666	21.03 ± 0.65	21.94 ± 0.35	20.21 ± 0.16	19.80 ± 0.11	19.14 ± 0.33	0.88 ± 0.08	4.86	26.31 ± 0.06	18.05
LFC 1494	587748929241612543	182.17507521	0.09046209	21.11 ± 0.47	20.55 ± 0.08	19.18 ± 0.04	18.51 ± 0.03	18.37 ± 0.11	1.28 ± 0.04	8.32	27.77 ± 0.12	8.55
LFC 1504	587748929241547195	181.98419233	0.08973886	23.10 ± 2.57	21.93 ± 0.26	20.63 ± 0.11	19.91 ± 0.09	20.17 ± 0.62	1.20 ± 0.08	6.52	26.93 ± 0.11	9.63
LFC 1511	587748929241547302	182.01602958	0.08979034	26.57 ± 1.53	22.25 ± 0.33	20.56 ± 0.11	20.17 ± 0.11	19.56 ± 0.31	0.93 ± 0.09	3.81	26.32 ± 0.05	23.49
LFC 1551	587748929241547404	182.04500046	0.09409627	20.81 ± 0.78	21.24 ± 0.32	20.14 ± 0.12	19.85 ± 0.20	18.59 ± 0.31	0.94 ± 0.18	3.81	25.60 ± 0.02	43.11
LFC 1585	587748929241612761	182.14289349	0.09547315	22.05 ± 0.81	21.58 ± 0.14	20.69 ± 0.09	20.09 ± 0.08	19.85 ± 0.40	0.82 ± 0.06	5.63	27.69 ± 0.16	6.40
LFC 1613	587748929241547377	182.03746603	0.09821770	21.68 ± 1.12	21.27 ± 0.21	20.67 ± 0.12	20.87 ± 0.33	20.90 ± 1.63	1.13 ± 0.25	5.36	28.04 ± 0.23	4.28
LFC 1628	587748929241612919	182.18963699	0.09910967	24.48 ± 5.72	21.73 ± 0.19	20.51 ± 0.09	19.86 ± 0.07	19.42 ± 0.24	0.97 ± 0.07	5.63	27.39 ± 0.18	5.64
LFC 1706	587748929241612709	182.12320511	0.10650842	24.18 ± 3.92	23.39 ± 0.76	20.93 ± 0.12	20.13 ± 0.09	20.30 ± 0.60	0.90 ± 0.07	5.63	26.75 ± 0.07	16.20
LFC 1739	587748929241612891	182.18384505	0.11087826	26.47 ± 1.33	22.02 ± 0.22	21.10 ± 0.15	20.35 ± 0.11	19.66 ± 0.30	0.86 ± 0.08	5.11	26.04 ± 0.06	16.67
LFC 1744	587748929241547049	182.06298264	0.11056218	21.95 ± 0.99	20.71 ± 0.09	19.81 ± 0.05	19.53 ± 0.08	18.83 ± 0.17	0.94 ± 0.05	3.81	25.83 ± 0.04	30.08
LFC 1772	587748929241547462	182.06821754	0.11339061	23.42 ± 3.70	20.71 ± 0.10	19.79 ± 0.05	19.59 ± 0.08	19.42 ± 0.32	1.12 ± 0.06	5.63	27.68 ± 0.18	5.43
LFC 1798	587748929241612610	182.16715984	0.11586897	21.00 ± 0.38	20.47 ± 0.06	19.98 ± 0.06	19.77 ± 0.08	19.56 ± 0.30	1.02 ± 0.06	5.36	28.49 ± 0.28	3.40
LFC 1802	587748929241612895	182.18515053	0.11669885	21.57 ± 0.67	20.50 ± 0.07	19.67 ± 0.05	19.36 ± 0.05	19.45 ± 0.28	1.08 ± 0.04	5.91	27.19 ± 0.12	8.38
LFC 1830	587748929241547045	182.04016040	0.12016220	21.76 ± 0.68	20.99 ± 0.09	19.96 ± 0.06	19.42 ± 0.05	19.14 ± 0.19	1.04 ± 0.05	5.91	27.23 ± 0.11	9.62
LFC 1833	587748929241547455	182.06517945	0.11903479	23.35 ± 2.54	22.03 ± 0.21	20.70 ± 0.10	20.35 ± 0.10	20.31 ± 0.48	0.77 ± 0.07	4.86	27.64 ± 0.11	9.29
LFC 1971	587748929241547674	182.02519253	0.13278430	22.90 ± 1.99	22.04 ± 0.28	21.12 ± 0.17	20.53 ± 0.15	19.57 ± 0.31	0.66 ± 0.09	5.63	28.08 ± 0.26	3.66
LFC 2007	587748929241547359	182.03063126	0.14040632	24.52 ± 7.77	21.46 ± 0.32	20.35 ± 0.25	19.83 ± 0.12	20.85 ± 2.04	0.83 ± 0.09	7.92	28.17 ± 0.17	5.92
LFC 2047	587748929241612936	182.19391201	0.14196037	22.08 ± 0.93	21.90 ± 0.23	20.46 ± 0.10	19.97 ± 0.08	19.54 ± 0.26	0.75 ± 0.05	5.11	27.76 ± 0.30	3.16
LFC 2106	587748929241612899	182.18552998	0.14664479	27.98 ± 1.83	21.63 ± 0.28	20.63 ± 0.13	20.10 ± 0.16	19.63 ± 0.52	1.33 ± 0.20	5.36	27.14 ± 0.14	7.49
LFC 2137	587748929241546927	182.04266233	0.14889641	20.96 ± 0.34	20.66 ± 0.07	20.24 ± 0.07	20.03 ± 0.07	20.16 ± 0.35	0.61 ± 0.04	3.81	26.65 ± 0.06	17.55
LFC 2159	587748929241547409	182.04708114	0.15187851	22.36 ± 1.68	21.56 ± 0.28	20.29 ± 0.09	19.78 ± 0.11	18.61 ± 0.18	0.99 ± 0.11	3.81	25.62 ± 0.03	37.61
LFC 2168	587748929241612454	182.16358958	0.15352032	21.00 ± 0.40	20.15 ± 0.05	19.54 ± 0.04	19.49 ± 0.06	19.31 ± 0.25	1.00 ± 0.05	3.81	25.96 ± 0.04	24.24
LFC 2177	587748929241547507	182.08226764	0.15471892	22.08 ± 1.34	20.84 ± 0.12	19.97 ± 0.07	19.96 ± 0.13	19.54 ± 0.39	1.08 ± 0.10	5.11	28.43 ± 0.23	4.22
LFC 2180	587748929241547188	181.97941963	0.15379342	21.97 ± 0.69	22.41 ± 0.29	21.77 ± 0.21	20.80 ± 0.13	19.94 ± 0.28	0.59 ± 0.06	3.59	26.97 ± 0.12	8.30
LFC 2181	587748929241612453	182.16386495	0.15506499	21.58 ± 0.57	20.25 ± 0.05	19.37 ± 0.03	19.02 ± 0.03	18.70 ± 0.12	0.86 ± 0.02	5.11	26.96 ± 0.09	11.76
LFC 2184	587748929241612341	182.11871678	0.16356295	18.04 ± 0.03	17.31 ± 0.01	17.11 ± 0.01	16.93 ± 0.01	16.88 ± 0.03	1.31 ± 0.01	11.15	25.75 ± 0.03	33.51
LFC 2310	587748929241547399	182.04442289	0.16755119	20.52 ± 0.43	20.35 ± 0.17	19.94 ± 0.55	19.79 ± 0.13	19.41 ± 0.44	0.96 ± 0.10	3.81	25.45 ± 0.03	32.83
LFC 2356	587748929241547486	182.07664357	0.17104016	22.10 ± 1.53	20.71 ± 0.12	20.30 ± 0.13	20.20 ± 0.17	19.85 ± 0.58	1.89 ± 0.25	3.81	25.13 ± 0.03	40.57
LFC 2362	587748929241547626	181.99416213	0.17046575	22.15 ± 1.73	22.37 ± 0.59	20.67 ± 0.12	19.92 ± 0.14	19.45 ± 0.44	1.31 ± 0.20	6.85	26.83 ± 0.11	9.03
LFC 2405	587748929241613208	182.15466470	0.17100824	26.50 ± 1.57	22.87 ± 0.58	21.57 ± 0.28	21.33 ± 0.32	20.20 ± 0.55	0.83 ± 0.22	3.46	28.01 ± 0.22	4.38
LFC 2414	587748929241546910	181.99490346	0.18396374	20.68 ± 0.38	19.61 ± 0.05	18.21 ± 0.03	17.72 ± 0.02	17.37 ± 0.06	1.35 ± 0.03	8.74	25.84 ± 0.04	27.76

Table 9—Continued

Our ID	SDSS ID	RA	Dec	u mag	g mag	r mag	i mag	z mag	R_{50} arcsec	sma arcsec	μ_i mag arcsec ⁻²	SNR
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
LFC 2422	587748929241612668	182.10793716	0.17687606	20.75 ± 0.49	19.90 ± 0.06	19.04 ± 0.03	18.60 ± 0.04	18.48 ± 0.18	1.82 ± 0.06	7.92	27.81 ± 0.13	7.88
LFC 2428	587748929241547039	182.03013212	0.17461684	25.92 ± 1.46	22.34 ± 0.19	21.19 ± 0.10	19.69 ± 0.04	18.87 ± 0.09	0.54 ± 0.02	5.36	26.70 ± 0.08	13.36
LFC 2472	587748929241612659	182.10234586	0.17914356	21.40 ± 0.55	21.13 ± 0.13	20.52 ± 0.11	20.15 ± 0.11	20.24 ± 0.57	1.41 ± 0.13	5.91	27.55 ± 0.22	4.38
LFC 2484	587748929241547440	182.05873627	0.17810661	20.71 ± 0.36	22.28 ± 0.40	20.69 ± 0.12	20.49 ± 0.17	20.73 ± 1.01	1.13 ± 0.14	4.41	28.26 ± 0.24	4.04
LFC 2503	587748929241547511	182.08428361	0.18036137	23.00 ± 1.28	22.84 ± 0.34	21.08 ± 0.11	20.27 ± 0.09	19.85 ± 0.23	0.83 ± 0.07	3.81	26.85 ± 0.08	13.58
LFC 2525	587748929241547315	182.01837362	0.18344027	22.20 ± 1.23	20.92 ± 0.11	20.26 ± 0.08	19.82 ± 0.09	20.03 ± 0.53	1.19 ± 0.09	3.81	26.06 ± 0.05	22.36
LFC 2539	587748929241547269	182.00438914	0.18432531	26.32 ± 1.18	22.34 ± 0.22	21.27 ± 0.13	20.45 ± 0.09	19.76 ± 0.22	0.81 ± 0.06	5.63	27.64 ± 0.15	6.55
LFC 2549	587748929241546928	182.04320837	0.18859279	20.21 ± 0.22	18.89 ± 0.02	18.09 ± 0.02	17.64 ± 0.01	17.40 ± 0.05	1.31 ± 0.02	9.63	27.12 ± 0.06	16.47
LFC 2568	587748929241612858	182.17630849	0.18495157	24.50 ± 4.10	21.48 ± 0.14	21.09 ± 0.16	20.41 ± 0.12	19.71 ± 0.29	1.06 ± 0.11	5.36	27.25 ± 0.11	9.23
LFC 2670	587748929241612781	182.14898514	0.19436667	23.89 ± 6.65	22.23 ± 0.44	20.71 ± 0.20	20.29 ± 0.17	19.74 ± 0.50	0.75 ± 0.11	3.81	26.77 ± 0.09	11.58
LFC 2692	587748929241547532	182.09162759	0.19664383	24.68 ± 4.77	21.14 ± 0.12	21.01 ± 0.17	21.05 ± 0.24	19.75 ± 0.36	1.00 ± 0.17	5.11	27.37 ± 0.24	4.08
LFC 2708	587748929241547280	182.00910743	0.19776933	26.73 ± 1.03	21.80 ± 0.16	20.52 ± 0.08	20.10 ± 0.08	21.97 ± 1.89	0.75 ± 0.05	3.81	26.39 ± 0.05	20.79
LFC 2709	587748929241612925	182.19123671	0.19795675	24.51 ± 3.69	22.14 ± 0.23	20.85 ± 0.12	20.18 ± 0.09	19.76 ± 0.28	0.78 ± 0.06	3.81	25.70 ± 0.05	22.75
LFC 2712	587748929241612471	182.18709793	0.20407222	21.26 ± 0.78	19.86 ± 0.08	18.79 ± 0.04	18.14 ± 0.05	17.60 ± 0.08	1.39 ± 0.08	7.55	25.39 ± 0.04	30.05
LFC 2719	587748929241612769	182.14348570	0.19877962	24.50 ± 4.44	22.34 ± 0.35	20.95 ± 0.17	20.52 ± 0.15	20.23 ± 0.50	0.95 ± 0.11	3.81	27.05 ± 0.11	9.11
LFC 2742	587748929241612470	182.18678466	0.20596149	19.62 ± 0.14	18.02 ± 0.01	17.17 ± 0.02	16.79 ± 0.02	16.45 ± 0.02	1.24 ± 0.03	11.15	25.05 ± 0.02	59.45
LFC 2782	588848900450418897	182.01556827	0.22315591	20.44 ± 0.26	19.21 ± 0.04	18.38 ± 0.03	17.87 ± 0.02	17.52 ± 0.08	2.01 ± 0.04	13.55	28.52 ± 0.20	4.99
LFC 2888	588848900450419376	182.06715352	0.21196476	23.59 ± 3.18	23.82 ± 1.29	21.21 ± 0.19	20.15 ± 0.13	19.29 ± 0.33	0.84 ± 0.11	3.81	26.25 ± 0.06	18.33
LFC 2955	588848900450484623	182.08566267	0.21663913	22.38 ± 0.65	21.54 ± 0.16	20.99 ± 0.12	20.47 ± 0.15	19.80 ± 0.28	0.82 ± 0.10	3.81	26.55 ± 0.05	20.26
LFC 2956	588848900450419362	182.06338063	0.21748981	23.25 ± 2.07	21.31 ± 0.16	20.57 ± 0.11	20.52 ± 0.19	19.32 ± 0.32	1.11 ± 0.14	3.81	26.46 ± 0.06	17.83
LFC 2988	588848900450484751	182.13663014	0.22157991	26.55 ± 1.12	22.48 ± 0.53	20.63 ± 0.15	19.70 ± 0.09	19.08 ± 0.28	1.27 ± 0.12	7.19	26.95 ± 0.08	13.64
LFC 3026	588848900450484758	182.13895296	0.22402556	26.28 ± 1.67	22.85 ± 0.59	21.00 ± 0.19	20.21 ± 0.19	19.70 ± 0.39	0.78 ± 0.15	5.36	27.44 ± 0.11	9.25
LFC 3072	588848900450484662	182.10294332	0.22805954	21.50 ± 0.39	22.40 ± 0.37	20.89 ± 0.15	20.51 ± 0.15	19.93 ± 0.41	0.97 ± 0.10	5.11	27.97 ± 0.22	4.54
LFC 3087	588848900450484864	182.17419997	0.22893448	23.65 ± 1.73	22.38 ± 0.26	21.13 ± 0.13	20.30 ± 0.09	19.55 ± 0.24	0.69 ± 0.05	4.41	27.32 ± 0.11	8.99
LFC 3094	588848900450418890	182.00116203	0.23227463	20.82 ± 0.26	20.15 ± 0.06	19.29 ± 0.04	19.05 ± 0.05	18.67 ± 0.17	1.24 ± 0.05	7.55	27.37 ± 0.10	10.48
LFC 3218	588848900450484539	182.09789162	0.23898822	21.86 ± 0.31	21.46 ± 0.09	20.78 ± 0.08	20.44 ± 0.09	20.29 ± 0.34	0.67 ± 0.04	4.63	27.67 ± 0.13	7.88
LFC 3274	588848900450484773	182.14357751	0.24490338	21.82 ± 0.47	21.58 ± 0.16	21.08 ± 0.16	20.56 ± 0.15	20.74 ± 0.80	1.36 ± 0.17	3.63	25.77 ± 0.11	9.55
LFC 3292	588848900450484557	182.15192415	0.24670492	21.31 ± 0.33	20.42 ± 0.06	19.98 ± 0.06	19.69 ± 0.07	19.74 ± 0.36	1.24 ± 0.10	6.21	26.98 ± 0.09	11.08
LFC 3329	588848900450484918	182.19474230	0.25043169	27.50 ± 0.44	22.27 ± 0.30	20.34 ± 0.08	19.68 ± 0.07	19.30 ± 0.21	0.86 ± 0.05	6.85	27.61 ± 0.17	5.88
LFC 3359	588848900450419242	182.01818845	0.25293238	21.98 ± 0.54	21.39 ± 0.14	20.70 ± 0.11	20.34 ± 0.12	19.74 ± 0.32	0.96 ± 0.08	4.41	27.50 ± 0.13	8.17
LFC 3381	588848900450484813	182.15539392	0.25472036	22.21 ± 0.60	22.16 ± 0.23	20.74 ± 0.10	19.97 ± 0.08	19.86 ± 0.31	0.98 ± 0.07	5.11	27.66 ± 0.18	5.51
LFC 3395	588848900450484512	182.15216700	0.25594856	22.38 ± 0.73	22.47 ± 0.28	20.83 ± 0.10	19.98 ± 0.07	19.37 ± 0.22	0.69 ± 0.04	5.36	28.11 ± 0.26	3.62
LFC 3423	588848900450419145	181.97726239	0.25980512	22.49 ± 0.94	22.29 ± 0.33	20.67 ± 0.11	20.10 ± 0.12	19.32 ± 0.28	0.99 ± 0.09	4.58	27.38 ± 0.08	12.80

Table 9—Continued

Our ID	SDSS ID	RA	Dec	u mag	g mag	r mag	i mag	z mag	R_{50} arcsec	sma arcsec	μ_i mag arcsec ⁻²	SNR
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
LFC 3427	588848900450485184	182.19056514	0.25979702	24.15 ± 4.28	22.69 ± 0.67	20.51 ± 0.16	20.05 ± 0.14	19.70 ± 0.47	1.25 ± 0.14	3.81	25.72 ± 0.05	23.09
LFC 3430	588848900450484421	182.18802200	0.26222895	20.05 ± 0.23	19.10 ± 0.05	18.31 ± 0.04	17.90 ± 0.03	17.53 ± 0.06	1.22 ± 0.05	11.15	27.05 ± 0.13	8.12
LFC 3458	588848900450484677	182.11331575	0.26184473	22.61 ± 0.83	22.15 ± 0.24	20.45 ± 0.08	19.96 ± 0.08	19.34 ± 0.20	0.87 ± 0.06	4.86	27.29 ± 0.10	9.98
LFC 3508	588848900450484456	182.11440196	0.26979325	22.33 ± 0.95	20.71 ± 0.09	19.20 ± 0.04	18.56 ± 0.04	18.29 ± 0.11	1.27 ± 0.06	8.74	26.75 ± 0.05	19.87
LFC 3562	588848900450484849	182.16773989	0.27028060	24.95 ± 2.49	22.83 ± 0.52	21.22 ± 0.18	20.56 ± 0.15	20.36 ± 0.56	0.70 ± 0.08	3.81	27.18 ± 0.10	10.64
LFC 3577	588848900450484730	182.13132095	0.27362923	21.71 ± 0.44	21.02 ± 0.10	19.82 ± 0.05	19.23 ± 0.05	18.76 ± 0.14	1.24 ± 0.04	6.52	27.66 ± 0.13	8.19
LFC 3632	588848900450484789	182.15159941	0.27678605	21.65 ± 0.37	22.64 ± 0.43	21.51 ± 0.21	20.49 ± 0.13	19.57 ± 0.26	0.82 ± 0.09	3.63	26.98 ± 0.12	8.88
LFC 3646	588848900450419263	182.02950009	0.28010104	25.32 ± 1.70	22.65 ± 0.34	20.81 ± 0.11	20.09 ± 0.10	19.60 ± 0.22	0.87 ± 0.08	3.81	25.80 ± 0.03	30.75
LFC 3760	588848900450484755	182.13792468	0.29146862	22.82 ± 1.56	21.28 ± 0.15	20.37 ± 0.09	19.68 ± 0.08	19.42 ± 0.30	1.39 ± 0.08	3.81	25.94 ± 0.04	29.63
LFC 3808	588848900450484737	182.13282007	0.29385752	21.87 ± 0.41	21.85 ± 0.17	20.66 ± 0.09	20.10 ± 0.08	19.49 ± 0.22	0.90 ± 0.05	4.63	27.63 ± 0.14	7.01
LFC 3878	588848900450484782	182.14952690	0.30063931	22.75 ± 1.14	21.68 ± 0.22	20.78 ± 0.12	20.36 ± 0.12	20.03 ± 0.41	0.99 ± 0.09	3.81	26.41 ± 0.10	10.57
LFC 3894	588848900450484724	182.12908347	0.30186681	22.38 ± 0.87	22.23 ± 0.30	20.51 ± 0.12	19.85 ± 0.10	19.28 ± 0.36	1.02 ± 0.10	3.81	26.25 ± 0.06	19.10
LFC 3912	588848900450485136	182.15368791	0.30241122	22.93 ± 1.52	24.90 ± 2.45	21.47 ± 0.25	20.38 ± 0.15	20.39 ± 0.70	0.95 ± 0.13	5.36	28.58 ± 0.27	3.51
LFC 3921	588848900450485183	182.18961582	0.30476227	21.93 ± 0.90	21.81 ± 0.34	20.92 ± 0.15	19.61 ± 0.11	19.52 ± 0.45	1.90 ± 0.31	4.63	26.06 ± 0.06	16.76
LFC 3927	588848900450418982	182.04376387	0.30459205	21.14 ± 0.28	20.34 ± 0.06	19.73 ± 0.05	19.61 ± 0.07	19.15 ± 0.21	1.06 ± 0.06	6.52	27.59 ± 0.12	8.80
LFC 3942	588848900450418913	182.06453615	0.30770545	26.56 ± 1.14	20.19 ± 0.09	19.56 ± 0.11	18.98 ± 0.09	18.38 ± 0.17	0.82 ± 0.07	7.92	28.04 ± 0.17	5.72
LFC 4009	588848900450484426	182.19366322	0.31301261	21.08 ± 0.24	19.97 ± 0.10	19.30 ± 0.12	18.83 ± 0.08	18.55 ± 0.25	0.97 ± 0.07	6.21	26.78 ± 0.21	4.68
LFC 4029	588848900450419357	182.06098194	0.31304151	23.92 ± 3.38	21.88 ± 0.29	20.52 ± 0.13	20.14 ± 0.14	19.47 ± 0.35	1.19 ± 0.14	5.36	27.21 ± 0.10	10.17
LFC 4059	588848900450484842	182.16621820	0.31743934	22.92 ± 1.86	20.77 ± 0.12	19.60 ± 0.07	19.17 ± 0.06	19.12 ± 0.28	1.52 ± 0.08	10.11	28.73 ± 0.29	3.31
LFC 4095	588848900450484850	182.16791279	0.32083467	24.49 ± 2.95	23.56 ± 0.86	21.29 ± 0.17	20.25 ± 0.10	19.53 ± 0.26	0.87 ± 0.07	4.20	26.97 ± 0.09	11.49
LFC 4188	588848900450484422	182.18935357	0.34142948	20.72 ± 0.21	19.48 ± 0.03	18.63 ± 0.02	18.30 ± 0.03	18.07 ± 0.09	1.10 ± 0.03	5.91	26.15 ± 0.07	16.01
LFC 4211	588848900450484464	182.14056980	0.33301123	24.20 ± 2.47	21.42 ± 0.13	19.73 ± 0.04	19.14 ± 0.04	18.66 ± 0.12	0.91 ± 0.04	5.63	27.11 ± 0.08	12.93
LFC 4248	588848900450484483	182.19749652	0.33861048	20.60 ± 0.30	19.58 ± 0.05	19.16 ± 0.05	18.95 ± 0.07	18.61 ± 0.23	1.56 ± 0.11	8.32	27.28 ± 0.25	3.82
LFC 4394	588848900450484756	182.13823066	0.36032260	22.99 ± 1.16	21.10 ± 0.09	20.57 ± 0.09	20.24 ± 0.10	19.91 ± 0.36	1.01 ± 0.08	3.81	26.79 ± 0.07	15.78
LFC 4400	588848900450419368	182.06512731	0.36449251	22.30 ± 1.01	22.11 ± 0.48	21.44 ± 0.24	20.40 ± 0.15	19.46 ± 0.36	1.25 ± 0.15	7.19	28.87 ± 0.20	5.03
LFC 4464	588848900450418988	182.06590839	0.35543977	21.96 ± 0.48	21.24 ± 0.10	20.40 ± 0.08	20.07 ± 0.09	19.28 ± 0.51	0.92 ± 0.07	5.63	26.95 ± 0.14	7.17
LFC 4488	588848900450484874	182.17876467	0.34574809	23.64 ± 2.27	22.28 ± 0.34	20.71 ± 0.13	19.85 ± 0.09	19.52 ± 0.29	1.09 ± 0.09	3.81	25.92 ± 0.04	26.16
LFC 4536	588848900450484565	182.18581747	0.33997004	21.40 ± 0.55	20.62 ± 0.11	20.27 ± 0.08	19.96 ± 0.15	19.76 ± 0.57	1.21 ± 0.20	3.81	25.96 ± 0.06	17.94
LFC 4654	588848900450484719	182.12587307	0.37281904	21.33 ± 0.40	21.32 ± 0.18	20.49 ± 0.12	20.50 ± 0.19	19.30 ± 0.29	1.05 ± 0.15	3.81	26.78 ± 0.12	8.38
LFC 4657	588848900450484500	182.08760528	0.34676179	23.68 ± 3.18	21.39 ± 0.20	19.76 ± 0.07	19.02 ± 0.06	18.73 ± 0.21	1.48 ± 0.11	7.92	27.80 ± 0.13	7.62
LFC 4681	588848900450484367	182.13718733	0.34356748	20.39 ± 0.16	19.49 ± 0.03	18.82 ± 0.03	18.51 ± 0.03	18.20 ± 0.10	1.24 ± 0.03	7.92	27.79 ± 0.13	7.58
LFC 4740	588848900450484476	182.18400953	0.34429933	21.98 ± 0.48	21.19 ± 0.11	19.85 ± 0.05	19.28 ± 0.05	18.90 ± 0.13	0.93 ± 0.05	6.52	26.80 ± 0.08	12.95
LFC 4763	588848900450484459	182.12762687	0.37994898	21.56 ± 0.25	21.03 ± 0.06	20.17 ± 0.05	19.66 ± 0.04	19.25 ± 0.14	0.69 ± 0.02	4.37	26.16 ± 0.09	11.10

Table 9—Continued

Our ID	SDSS ID	RA	Dec	u	g	r	i	z	R_{50}	sma	μ_i	SNR
				mag	mag	mag	mag	mag	arcsec	arcsec	mag arcsec ⁻²	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)

Note. — Column (1) gives ID of the sample galaxies in SExtractor catalog generated by us. Quantities from column (2) to column (10) are obtained from the PhotoObjAll table of the SDSS DR7 photometric catalog (Abazajian et al. 2009). Column (2) gives a unique SDSS identifier which is listed as objID in the PhotoObjAll table Columns (3) and (4) give J2000 right ascension and declination in degree. Columns (5) to (9) give extinction corrected Petrosian magnitudes in u , g , r , i and z bands. Column (10) gives Petrosian half-light radius in a arcsec. Columns (11) to (13) give length of semi-major axis (in arcsec), i -band surface brightness (in $mag/arcsec^{-2}$) and signal to noise ratio respectively of the outermost isophote in which ellipse can be fitted (see Section 4).

Table 10. Derived parametrs for sample galaxies

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 462	0.89
		$3.52\text{e-}03 \pm 2.25\text{e-}03$	$-1.25\text{e-}03 \pm 2.16\text{e-}03$	0.057 ± 0.0058	60.37 ± 3.02
		$7.01\text{e-}03 \pm 4.73\text{e-}03$	$6.91\text{e-}03 \pm 3.09\text{e-}03$	0.048 ± 0.0063	1.87 ± 3.67
	
LFC 525	0.97
		$5.52\text{e-}03 \pm 1.89\text{e-}03$	$-1.09\text{e-}03 \pm 1.53\text{e-}03$	0.070 ± 0.0047	8.47 ± 2.02
		$-6.61\text{e-}03 \pm 5.70\text{e-}03$	$1.21\text{e-}02 \pm 5.41\text{e-}03$	0.023 ± 0.0117	8.22 ± 11.97
	
LFC 554	0.65	$-4.29\text{e-}03 \pm 1.96\text{e-}03$	$1.00\text{e-}02 \pm 1.82\text{e-}03$	0.094 ± 0.0040	31.47 ± 1.26
		$-8.35\text{e-}03 \pm 3.77\text{e-}03$	$3.49\text{e-}03 \pm 3.75\text{e-}03$	0.115 ± 0.0071	11.55 ± 1.92
	
	
LFC 558	0.68	$6.77\text{e-}03 \pm 1.47\text{e-}03$	$1.88\text{e-}03 \pm 1.42\text{e-}03$	0.038 ± 0.0035	1.71 ± 2.62
		$4.92\text{e-}03 \pm 1.60\text{e-}03$	$1.12\text{e-}04 \pm 1.56\text{e-}03$	0.081 ± 0.0036	5.48 ± 1.19
		$-1.10\text{e-}02 \pm 1.19\text{e-}02$	$-4.33\text{e-}03 \pm 1.28\text{e-}02$	0.164 ± 0.0248	119.00 ± 4.33
	
LFC 622	0.59	$-4.88\text{e-}04 \pm 1.06\text{e-}03$	$1.51\text{e-}02 \pm 4.42\text{e-}04$	0.358 ± 0.0018	1.64 ± 0.18
		$-1.18\text{e-}03 \pm 8.45\text{e-}04$	$1.29\text{e-}02 \pm 4.69\text{e-}04$	0.421 ± 0.0013	0.78 ± 0.12
		$4.78\text{e-}02 \pm 4.35\text{e-}03$	$-1.47\text{e-}02 \pm 3.35\text{e-}03$	0.310 ± 0.0075	0.89 ± 0.87
		$3.79\text{e-}02 \pm 1.57\text{e-}02$	$-2.56\text{e-}02 \pm 1.45\text{e-}02$	0.259 ± 0.0257	17.35 ± 3.29
LFC 633	0.66	$5.30\text{e-}03 \pm 7.48\text{e-}04$	$-2.57\text{e-}03 \pm 6.74\text{e-}04$	0.213 ± 0.0016	0.83 ± 0.24
		$1.59\text{e-}02 \pm 1.24\text{e-}03$	$-3.83\text{e-}03 \pm 1.11\text{e-}03$	0.219 ± 0.0027	6.21 ± 0.40
		$1.64\text{e-}02 \pm 4.29\text{e-}03$	$-6.56\text{e-}04 \pm 4.22\text{e-}03$	0.269 ± 0.0074	26.91 ± 0.91
		$-2.51\text{e-}01 \pm 7.91\text{e-}02$	$-2.39\text{e-}02 \pm 2.74\text{e-}02$	0.102 ± 0.0591	0.00 ± 16.88
LFC 663	0.74	$-6.50\text{e-}03 \pm 1.38\text{e-}03$	$-6.86\text{e-}03 \pm 1.21\text{e-}03$	0.284 ± 0.0028	0.22 ± 0.33
		$-9.27\text{e-}03 \pm 1.63\text{e-}03$	$-1.92\text{e-}03 \pm 1.56\text{e-}03$	0.282 ± 0.0031	1.68 ± 0.38
	
	
LFC 677	0.92	$5.66\text{e-}03 \pm 3.87\text{e-}03$	$-7.97\text{e-}03 \pm 3.12\text{e-}03$	0.474 ± 0.0060	0.68 ± 0.51
		$6.06\text{e-}03 \pm 2.19\text{e-}03$	$6.94\text{e-}04 \pm 1.89\text{e-}03$	0.443 ± 0.0036	2.67 ± 0.31
		$1.20\text{e-}03 \pm 9.19\text{e-}03$	$-3.97\text{e-}02 \pm 9.54\text{e-}03$	0.460 ± 0.0146	1.74 ± 1.24
	
LFC 725	0.74	$1.98\text{e-}02 \pm 4.50\text{e-}03$	$4.01\text{e-}03 \pm 3.26\text{e-}03$	0.084 ± 0.0102	19.88 ± 3.43
		$1.76\text{e-}02 \pm 3.79\text{e-}03$	$1.26\text{e-}02 \pm 3.55\text{e-}03$	0.095 ± 0.0083	18.92 ± 2.21
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a^3}{a} \rangle$ (3)	$\langle \frac{a^4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 741	0.55	$-7.78\text{e-}03 \pm 3.24\text{e-}03$	$4.65\text{e-}03 \pm 3.10\text{e-}03$	0.218 ± 0.0073	0.94 ± 1.10
		$-5.23\text{e-}03 \pm 2.10\text{e-}03$	$-2.21\text{e-}03 \pm 1.97\text{e-}03$	0.213 ± 0.0042	13.57 ± 0.64
		$-5.80\text{e-}02 \pm 2.05\text{e-}02$	$-2.09\text{e-}02 \pm 1.57\text{e-}02$	0.307 ± 0.0310	13.76 ± 3.25
	
LFC 824	0.83	$-1.15\text{e-}03 \pm 2.14\text{e-}03$	$-4.45\text{e-}03 \pm 1.69\text{e-}03$	0.059 ± 0.0045	0.67 ± 2.27
		$-6.73\text{e-}03 \pm 1.89\text{e-}03$	$2.00\text{e-}03 \pm 1.43\text{e-}03$	0.053 ± 0.0040	23.31 ± 2.03
		$-1.03\text{e-}02 \pm 8.73\text{e-}03$	$-9.41\text{e-}03 \pm 7.96\text{e-}03$	0.148 ± 0.0062	0.00 ± 1.30
	
LFC 848	0.65	$7.87\text{e-}03 \pm 1.78\text{e-}03$	$2.35\text{e-}03 \pm 1.73\text{e-}03$	0.042 ± 0.0038	8.66 ± 2.63
		$2.55\text{e-}03 \pm 3.05\text{e-}03$	$3.14\text{e-}03 \pm 2.82\text{e-}03$	0.054 ± 0.0064	23.19 ± 3.21
	
	
LFC 925	0.47	$2.51\text{e-}03 \pm 1.37\text{e-}03$	$3.86\text{e-}03 \pm 1.31\text{e-}03$	0.183 ± 0.0029	0.89 ± 0.51
		$3.34\text{e-}03 \pm 1.48\text{e-}03$	$-3.67\text{e-}03 \pm 1.43\text{e-}03$	0.166 ± 0.0032	0.38 ± 0.61
	
	
LFC 954	0.59	$-1.10\text{e-}02 \pm 1.53\text{e-}03$	$5.62\text{e-}03 \pm 1.14\text{e-}03$	0.152 ± 0.0034	0.92 ± 0.83
		$-1.02\text{e-}04 \pm 5.82\text{e-}04$	$1.73\text{e-}04 \pm 5.41\text{e-}04$	0.129 ± 0.0041	0.00 ± 3.35
	
	
LFC 985	0.71	$-9.05\text{e-}03 \pm 1.46\text{e-}03$	$4.85\text{e-}03 \pm 1.31\text{e-}03$	0.092 ± 0.0036	18.64 ± 1.14
		$-2.43\text{e-}02 \pm 2.81\text{e-}03$	$1.56\text{e-}02 \pm 2.57\text{e-}03$	0.147 ± 0.0057	12.39 ± 1.20
		$1.56\text{e-}02 \pm 1.48\text{e-}02$	$1.58\text{e-}02 \pm 1.53\text{e-}02$	0.156 ± 0.0284	22.63 ± 5.13
	
LFC 988	0.40	$-8.92\text{e-}04 \pm 1.96\text{e-}03$	$1.06\text{e-}03 \pm 1.91\text{e-}03$	0.088 ± 0.0040	9.96 ± 1.39
		$1.56\text{e-}02 \pm 3.23\text{e-}03$	$3.03\text{e-}03 \pm 3.06\text{e-}03$	0.074 ± 0.0067	26.19 ± 2.50
	
	
LFC 1013	0.53	$8.08\text{e-}03 \pm 1.27\text{e-}03$	$5.05\text{e-}04 \pm 1.05\text{e-}03$	0.522 ± 0.0020	0.63 ± 0.16
		$1.19\text{e-}02 \pm 1.81\text{e-}03$	$5.15\text{e-}03 \pm 1.53\text{e-}03$	0.515 ± 0.0023	0.16 ± 0.19
	
	
LFC 1039	0.67	$3.23\text{e-}04 \pm 2.10\text{e-}03$	$6.03\text{e-}03 \pm 1.97\text{e-}03$	0.420 ± 0.0035	1.42 ± 0.32
		$-4.47\text{e-}03 \pm 1.85\text{e-}03$	$1.05\text{e-}02 \pm 1.72\text{e-}03$	0.409 ± 0.0032	0.56 ± 0.29
		$-4.50\text{e-}03 \pm 1.44\text{e-}02$	$-2.66\text{e-}02 \pm 1.42\text{e-}02$	0.329 ± 0.0227	1.08 ± 2.61
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 1042	0.61	$-4.99\text{e-}03 \pm 2.21\text{e-}03$	$9.33\text{e-}03 \pm 1.64\text{e-}03$	0.244 ± 0.0045	0.64 ± 0.61
		$-1.63\text{e-}02 \pm 1.87\text{e-}03$	$4.54\text{e-}03 \pm 1.67\text{e-}03$	0.248 ± 0.0040	12.68 ± 0.54
		$-1.41\text{e-}02 \pm 1.23\text{e-}02$	$-3.13\text{e-}02 \pm 1.04\text{e-}02$	0.193 ± 0.0211	27.04 ± 3.03
	
LFC 1044	0.78	$-4.15\text{e-}03 \pm 5.67\text{e-}04$	$2.31\text{e-}03 \pm 5.44\text{e-}04$	0.389 ± 0.0011	0.28 ± 0.11
		$-5.92\text{e-}03 \pm 8.77\text{e-}04$	$-6.89\text{e-}04 \pm 8.53\text{e-}04$	0.365 ± 0.0016	0.80 ± 0.16
		$4.11\text{e-}02 \pm 8.83\text{e-}03$	$-7.08\text{e-}02 \pm 9.41\text{e-}03$	0.267 ± 0.0138	10.98 ± 1.77
		$-6.87\text{e-}03 \pm 1.93\text{e-}02$	$-1.56\text{e-}02 \pm 1.96\text{e-}02$	0.194 ± 0.0354	4.73 ± 5.91
LFC 1056	0.83	$3.53\text{e-}03 \pm 8.96\text{e-}04$	$3.99\text{e-}03 \pm 7.94\text{e-}04$	0.387 ± 0.0017	1.04 ± 0.16
		$2.21\text{e-}03 \pm 8.22\text{e-}04$	$7.50\text{e-}03 \pm 6.95\text{e-}04$	0.383 ± 0.0014	0.76 ± 0.13
		$-2.18\text{e-}02 \pm 4.81\text{e-}03$	$3.87\text{e-}02 \pm 4.72\text{e-}03$	0.347 ± 0.0078	5.04 ± 0.79
		$-8.47\text{e-}03 \pm 9.42\text{e-}03$	$1.09\text{e-}02 \pm 9.73\text{e-}03$	0.377 ± 0.0147	2.68 ± 1.41
LFC 1091	0.65
		$-1.57\text{e-}02 \pm 2.23\text{e-}03$	$2.82\text{e-}03 \pm 1.91\text{e-}03$	0.151 ± 0.0048	23.31 ± 1.01
		$4.82\text{e-}02 \pm 7.96\text{e-}03$	$2.12\text{e-}02 \pm 6.42\text{e-}03$	0.162 ± 0.0158	39.45 ± 3.04
		$3.16\text{e-}02 \pm 2.08\text{e-}02$	$5.66\text{e-}02 \pm 2.07\text{e-}02$	0.166 ± 0.0348	3.53 ± 6.58
LFC 1124	0.60	$1.45\text{e-}02 \pm 1.43\text{e-}03$	$-4.58\text{e-}03 \pm 1.33\text{e-}03$	0.196 ± 0.0040	2.09 ± 0.66
		$1.95\text{e-}03 \pm 2.58\text{e-}03$	$4.26\text{e-}03 \pm 2.47\text{e-}03$	0.162 ± 0.0053	2.83 ± 1.02
	
	
LFC 1204	0.62	$-7.90\text{e-}03 \pm 1.49\text{e-}03$	$2.78\text{e-}04 \pm 1.44\text{e-}03$	0.319 ± 0.0029	1.52 ± 0.32
		$-1.31\text{e-}02 \pm 2.12\text{e-}03$	$1.26\text{e-}03 \pm 2.02\text{e-}03$	0.319 ± 0.0040	6.17 ± 0.43
		$6.04\text{e-}04 \pm 1.60\text{e-}02$	$1.79\text{e-}02 \pm 1.69\text{e-}02$	0.149 ± 0.0309	92.06 ± 6.51
	
LFC 1239	0.71	$-1.38\text{e-}03 \pm 1.77\text{e-}03$	$8.07\text{e-}03 \pm 1.32\text{e-}03$	0.207 ± 0.0034	0.62 ± 0.53
		$2.86\text{e-}04 \pm 1.03\text{e-}03$	$1.02\text{e-}02 \pm 8.02\text{e-}04$	0.196 ± 0.0020	9.01 ± 0.34
		$3.05\text{e-}02 \pm 5.35\text{e-}03$	$-2.01\text{e-}03 \pm 4.76\text{e-}03$	0.213 ± 0.0097	7.93 ± 1.43
		$2.91\text{e-}02 \pm 1.43\text{e-}02$	$-3.22\text{e-}02 \pm 1.32\text{e-}02$	0.256 ± 0.0233	7.56 ± 3.18
LFC 1263	0.46	$-1.68\text{e-}03 \pm 3.48\text{e-}03$	$-1.21\text{e-}02 \pm 3.40\text{e-}03$	0.110 ± 0.0078	60.72 ± 2.38
	
	
	
LFC 1273	0.48	$-8.47\text{e-}03 \pm 2.83\text{e-}03$	$-1.04\text{e-}02 \pm 2.63\text{e-}03$	0.124 ± 0.0057	5.37 ± 1.39
		$4.56\text{e-}03 \pm 4.35\text{e-}03$	$-1.69\text{e-}02 \pm 4.21\text{e-}03$	0.137 ± 0.0083	2.80 ± 1.85
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 1308	0.90	$-5.60\text{e-}03 \pm 2.34\text{e-}03$	$-1.20\text{e-}02 \pm 2.16\text{e-}03$	0.127 ± 0.0048	2.76 ± 1.18
		$-9.58\text{e-}03 \pm 2.45\text{e-}03$	$-1.72\text{e-}02 \pm 2.21\text{e-}03$	0.120 ± 0.0051	15.51 ± 1.35
		$1.16\text{e-}01 \pm 1.29\text{e-}01$	$4.77\text{e-}02 \pm 8.49\text{e-}02$	0.098 ± 0.1470	0.00 ± 45.16
	
LFC 1369	0.69	$-4.75\text{e-}03 \pm 1.83\text{e-}03$	$4.57\text{e-}04 \pm 1.75\text{e-}03$	0.191 ± 0.0038	7.31 ± 0.64
		$-1.39\text{e-}02 \pm 3.08\text{e-}03$	$4.81\text{e-}03 \pm 2.63\text{e-}03$	0.199 ± 0.0061	5.73 ± 1.00
	
	
LFC 1385	0.69	$-1.76\text{e-}03 \pm 1.59\text{e-}03$	$3.06\text{e-}03 \pm 1.49\text{e-}03$	0.041 ± 0.0033	19.91 ± 2.40
		$2.03\text{e-}03 \pm 2.01\text{e-}03$	$-8.56\text{e-}04 \pm 1.91\text{e-}03$	0.051 ± 0.0039	38.83 ± 1.91
	
	
LFC 1411	0.71	$-1.17\text{e-}02 \pm 2.64\text{e-}03$	$6.91\text{e-}03 \pm 2.22\text{e-}03$	0.308 ± 0.0051	0.87 ± 0.57
		$-1.07\text{e-}02 \pm 3.19\text{e-}03$	$-7.02\text{e-}03 \pm 2.83\text{e-}03$	0.269 ± 0.0059	45.80 ± 0.75
	
	
LFC 1437	0.95	$1.87\text{e-}03 \pm 2.50\text{e-}03$	$2.37\text{e-}03 \pm 2.44\text{e-}03$	0.296 ± 0.0045	1.10 ± 0.53
		$1.57\text{e-}04 \pm 2.20\text{e-}03$	$1.59\text{e-}02 \pm 1.95\text{e-}03$	0.292 ± 0.0039	1.43 ± 0.46
	
	
LFC 1456	0.83	$-6.91\text{e-}03 \pm 1.69\text{e-}03$	$5.56\text{e-}03 \pm 1.50\text{e-}03$	0.208 ± 0.0036	3.23 ± 0.56
		$1.06\text{e-}03 \pm 1.92\text{e-}03$	$-7.96\text{e-}04 \pm 1.69\text{e-}03$	0.201 ± 0.0041	5.96 ± 0.66
	
	
LFC 1459	0.78	$1.74\text{e-}02 \pm 2.78\text{e-}03$	$1.13\text{e-}02 \pm 2.50\text{e-}03$	0.078 ± 0.0066	3.23 ± 2.55
		$7.29\text{e-}03 \pm 1.77\text{e-}03$	$1.70\text{e-}03 \pm 1.63\text{e-}03$	0.079 ± 0.0038	56.75 ± 1.47
		$1.17\text{e-}02 \pm 1.29\text{e-}02$	$-2.51\text{e-}02 \pm 1.26\text{e-}02$	0.134 ± 0.0245	15.75 ± 5.22
	
LFC 1477	0.66	$-5.69\text{e-}03 \pm 1.41\text{e-}03$	$1.43\text{e-}03 \pm 1.38\text{e-}03$	0.102 ± 0.0030	6.13 ± 0.89
		$-9.68\text{e-}03 \pm 2.13\text{e-}03$	$1.73\text{e-}03 \pm 2.03\text{e-}03$	0.138 ± 0.0049	73.88 ± 1.07
		$3.02\text{e-}02 \pm 1.87\text{e-}02$	$1.99\text{e-}02 \pm 1.82\text{e-}02$	0.096 ± 0.0375	19.89 ± 11.31
	
LFC 1494	0.88	$-3.00\text{e-}03 \pm 7.61\text{e-}04$	$2.66\text{e-}03 \pm 7.04\text{e-}04$	0.099 ± 0.0016	0.26 ± 0.50
		$4.54\text{e-}03 \pm 1.19\text{e-}03$	$-1.21\text{e-}03 \pm 1.16\text{e-}03$	0.126 ± 0.0023	9.31 ± 0.56
		$3.10\text{e-}03 \pm 4.96\text{e-}03$	$-1.89\text{e-}02 \pm 5.02\text{e-}03$	0.187 ± 0.0091	32.22 ± 1.56
		$-9.84\text{e-}02 \pm 5.20\text{e-}02$	$-2.21\text{e-}02 \pm 3.58\text{e-}02$	0.139 ± 0.0687	0.00 ± 15.20

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 1504	0.63	$-7.38\text{e-}03 \pm 2.23\text{e-}03$	$2.73\text{e-}03 \pm 2.03\text{e-}03$	0.097 ± 0.0046	1.52 ± 1.39
		$1.05\text{e-}03 \pm 3.57\text{e-}03$	$4.79\text{e-}03 \pm 3.43\text{e-}03$	0.152 ± 0.0069	40.98 ± 1.45
		$5.02\text{e-}03 \pm 1.28\text{e-}02$	$4.83\text{e-}02 \pm 1.39\text{e-}02$	0.343 ± 0.0239	0.94 ± 2.04
	
LFC 1511	0.50	$-8.59\text{e-}03 \pm 1.68\text{e-}03$	$7.32\text{e-}03 \pm 1.49\text{e-}03$	0.167 ± 0.0035	5.04 ± 0.67
		$-6.04\text{e-}03 \pm 2.34\text{e-}03$	$-4.44\text{e-}03 \pm 2.26\text{e-}03$	0.187 ± 0.0045	4.94 ± 0.76
	
	
LFC 1551	0.91	$-5.49\text{e-}03 \pm 1.43\text{e-}03$	$-5.26\text{e-}03 \pm 1.31\text{e-}03$	0.061 ± 0.0033	1.54 ± 1.61
		$7.59\text{e-}03 \pm 2.77\text{e-}03$	$1.15\text{e-}02 \pm 2.60\text{e-}03$	0.068 ± 0.0054	38.81 ± 2.13
	
	
LFC 1585	0.67	$-7.80\text{e-}03 \pm 3.30\text{e-}03$	$7.22\text{e-}03 \pm 2.81\text{e-}03$	0.108 ± 0.0073	0.00 ± 2.11
		$-6.85\text{e-}03 \pm 1.55\text{e-}03$	$7.01\text{e-}03 \pm 1.32\text{e-}03$	0.106 ± 0.0033	1.34 ± 0.96
		$-1.67\text{e-}02 \pm 7.70\text{e-}03$	$-1.72\text{e-}02 \pm 7.23\text{e-}03$	0.131 ± 0.0146	61.31 ± 3.58
		$2.16\text{e-}02 \pm 2.29\text{e-}02$	$-2.72\text{e-}02 \pm 2.27\text{e-}02$	0.197 ± 0.0406	29.55 ± 6.40
LFC 1613	0.71	$-2.10\text{e-}02 \pm 2.67\text{e-}03$	$-3.34\text{e-}04 \pm 2.47\text{e-}03$	0.144 ± 0.0059	23.21 ± 1.32
		$-3.74\text{e-}02 \pm 6.94\text{e-}03$	$2.60\text{e-}02 \pm 5.64\text{e-}03$	0.112 ± 0.0128	5.01 ± 3.45
	
	
LFC 1628	0.65	$-1.49\text{e-}02 \pm 2.68\text{e-}03$	$-5.31\text{e-}04 \pm 2.21\text{e-}03$	0.132 ± 0.0061	0.01 ± 1.45
		$2.58\text{e-}03 \pm 2.25\text{e-}03$	$4.51\text{e-}03 \pm 2.15\text{e-}03$	0.107 ± 0.0045	5.99 ± 1.30
		$-8.63\text{e-}03 \pm 1.26\text{e-}02$	$-2.63\text{e-}03 \pm 1.18\text{e-}02$	0.115 ± 0.0223	22.58 ± 5.52
		$-2.69\text{e-}02 \pm 6.40\text{e-}02$	$-5.37\text{e-}02 \pm 6.93\text{e-}02$	0.131 ± 0.1177	0.00 ± 27.61
LFC 1706	0.84	$-5.04\text{e-}03 \pm 2.34\text{e-}03$	$-4.92\text{e-}03 \pm 2.18\text{e-}03$	0.071 ± 0.0058	0.14 ± 2.51
		$-1.52\text{e-}02 \pm 2.07\text{e-}03$	$-2.64\text{e-}03 \pm 1.96\text{e-}03$	0.056 ± 0.0045	1.81 ± 2.35
		$2.01\text{e-}02 \pm 7.47\text{e-}03$	$4.94\text{e-}03 \pm 6.35\text{e-}03$	0.146 ± 0.0137	19.54 ± 2.38
		$1.27\text{e-}02 \pm 2.33\text{e-}02$	$2.30\text{e-}02 \pm 2.59\text{e-}02$	0.264 ± 0.0407	0.00 ± 4.54
LFC 1739	0.65	$-1.06\text{e-}02 \pm 2.97\text{e-}03$	$-6.39\text{e-}04 \pm 2.62\text{e-}03$	0.041 ± 0.0063	16.89 ± 4.00
		$-8.15\text{e-}03 \pm 3.93\text{e-}03$	$-8.50\text{e-}03 \pm 3.92\text{e-}03$	0.142 ± 0.0079	10.15 ± 1.43
		$-1.57\text{e-}02 \pm 1.34\text{e-}02$	$-4.46\text{e-}02 \pm 1.43\text{e-}02$	0.331 ± 0.0248	2.96 ± 2.80
	
LFC 1744	0.86	$5.52\text{e-}03 \pm 1.02\text{e-}03$	$5.05\text{e-}03 \pm 7.75\text{e-}04$	0.301 ± 0.0023	0.22 ± 0.26
		$3.81\text{e-}03 \pm 8.59\text{e-}04$	$3.00\text{e-}03 \pm 8.02\text{e-}04$	0.287 ± 0.0019	1.14 ± 0.23
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 1772	0.57	$5.83\text{e-}04 \pm 9.35\text{e-}04$	$5.43\text{e-}04 \pm 8.97\text{e-}04$	0.205 ± 0.0019	0.82 ± 0.29
		$1.74\text{e-}04 \pm 1.24\text{e-}03$	$6.89\text{e-}04 \pm 1.19\text{e-}03$	0.206 ± 0.0024	0.73 ± 0.37
		$-9.21\text{e-}03 \pm 1.05\text{e-}02$	$2.95\text{e-}02 \pm 1.08\text{e-}02$	0.216 ± 0.0187	21.70 ± 2.87
	
LFC 1798	0.49	$-8.95\text{e-}03 \pm 1.55\text{e-}03$	$3.49\text{e-}03 \pm 1.43\text{e-}03$	0.151 ± 0.0036	0.99 ± 0.75
		$-3.35\text{e-}03 \pm 1.38\text{e-}03$	$5.89\text{e-}03 \pm 1.29\text{e-}03$	0.138 ± 0.0029	1.10 ± 0.66
		$-8.38\text{e-}03 \pm 9.41\text{e-}03$	$-5.75\text{e-}03 \pm 9.16\text{e-}03$	0.103 ± 0.0182	17.80 ± 5.41
	
LFC 1802	0.88	$-6.10\text{e-}03 \pm 1.16\text{e-}03$	$1.12\text{e-}03 \pm 9.98\text{e-}04$	0.132 ± 0.0029	1.05 ± 0.69
		$8.35\text{e-}03 \pm 1.85\text{e-}03$	$-1.95\text{e-}03 \pm 1.78\text{e-}03$	0.137 ± 0.0044	16.99 ± 0.99
		$-6.41\text{e-}02 \pm 2.00\text{e-}02$	$1.12\text{e-}02 \pm 1.38\text{e-}02$	0.147 ± 0.0299	25.61 ± 5.68
	
LFC 1830	0.52	$-3.34\text{e-}03 \pm 1.10\text{e-}03$	$1.45\text{e-}03 \pm 1.08\text{e-}03$	0.152 ± 0.0024	1.01 ± 0.49
		$-1.73\text{e-}03 \pm 1.09\text{e-}03$	$4.14\text{e-}03 \pm 1.03\text{e-}03$	0.134 ± 0.0022	16.08 ± 0.53
		$-1.72\text{e-}02 \pm 1.10\text{e-}02$	$-2.76\text{e-}02 \pm 1.04\text{e-}02$	0.052 ± 0.0201	27.99 ± 9.29
		$-4.44\text{e-}02 \pm 3.77\text{e-}02$	$-1.00\text{e-}01 \pm 4.78\text{e-}02$	0.020 ± 0.0706	0.00 ± 100.30
LFC 1833	0.46	$-1.11\text{e-}03 \pm 2.27\text{e-}03$	$-1.56\text{e-}03 \pm 2.13\text{e-}03$	0.085 ± 0.0048	6.50 ± 1.73
		$4.06\text{e-}03 \pm 3.05\text{e-}03$	$2.56\text{e-}03 \pm 2.95\text{e-}03$	0.048 ± 0.0064	61.70 ± 3.57
		$4.73\text{e-}03 \pm 1.97\text{e-}02$	$5.88\text{e-}03 \pm 1.68\text{e-}02$	0.125 ± 0.0338	5.59 ± 8.07
	
LFC 1971	0.77	$-2.49\text{e-}03 \pm 3.12\text{e-}03$	$1.54\text{e-}03 \pm 2.82\text{e-}03$	0.088 ± 0.0073	4.86 ± 2.51
		$-2.53\text{e-}02 \pm 2.81\text{e-}03$	$1.52\text{e-}02 \pm 2.55\text{e-}03$	0.130 ± 0.0064	11.75 ± 1.35
		$4.12\text{e-}02 \pm 1.11\text{e-}02$	$5.66\text{e-}03 \pm 9.55\text{e-}03$	0.149 ± 0.0186	1.84 ± 3.90
	
LFC 2007	0.79	$1.58\text{e-}03 \pm 1.30\text{e-}03$	$3.28\text{e-}03 \pm 1.23\text{e-}03$	0.116 ± 0.0026	4.08 ± 0.69
		$1.87\text{e-}02 \pm 2.59\text{e-}03$	$-1.54\text{e-}02 \pm 2.43\text{e-}03$	0.157 ± 0.0054	3.36 ± 1.00
		$1.63\text{e-}03 \pm 9.97\text{e-}03$	$-2.98\text{e-}02 \pm 1.01\text{e-}02$	0.295 ± 0.0160	3.59 ± 2.04
	
LFC 2047	0.67	$-5.40\text{e-}03 \pm 2.31\text{e-}03$	$2.51\text{e-}03 \pm 2.27\text{e-}03$	0.176 ± 0.0045	2.23 ± 0.81
		$-1.60\text{e-}03 \pm 3.16\text{e-}03$	$2.49\text{e-}03 \pm 3.11\text{e-}03$	0.197 ± 0.0060	49.67 ± 0.98
		$2.80\text{e-}02 \pm 1.73\text{e-}02$	$-3.66\text{e-}02 \pm 1.63\text{e-}02$	0.214 ± 0.0348	35.92 ± 4.53
	
LFC 2106	0.94	$2.73\text{e-}02 \pm 2.46\text{e-}03$	$9.86\text{e-}03 \pm 2.02\text{e-}03$	0.389 ± 0.0046	1.73 ± 0.43
		$7.31\text{e-}03 \pm 3.03\text{e-}03$	$1.08\text{e-}02 \pm 2.90\text{e-}03$	0.393 ± 0.0048	4.05 ± 0.46
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 2137	0.85	$5.57\text{e-}02 \pm 5.20\text{e-}03$	$2.11\text{e-}02 \pm 2.78\text{e-}03$	0.130 ± 0.0134	1.52 ± 3.15
		$6.02\text{e-}02 \pm 3.61\text{e-}03$	$2.11\text{e-}02 \pm 2.01\text{e-}03$	0.190 ± 0.0072	16.65 ± 1.21
	
	
LFC 2159	0.65	$2.05\text{e-}03 \pm 1.51\text{e-}03$	$-1.78\text{e-}03 \pm 1.29\text{e-}03$	0.092 ± 0.0031	9.20 ± 1.02
		$3.83\text{e-}03 \pm 1.67\text{e-}03$	$1.50\text{e-}03 \pm 1.61\text{e-}03$	0.143 ± 0.0031	6.13 ± 0.61
	
	
LFC 2168	0.67	$7.89\text{e-}03 \pm 1.09\text{e-}03$	$-2.46\text{e-}03 \pm 1.02\text{e-}03$	0.183 ± 0.0028	0.46 ± 0.50
		$1.02\text{e-}02 \pm 1.51\text{e-}03$	$-2.65\text{e-}03 \pm 1.45\text{e-}03$	0.145 ± 0.0034	25.89 ± 0.76
	
	
LFC 2177	0.60	$5.98\text{e-}03 \pm 1.41\text{e-}03$	$-1.17\text{e-}03 \pm 1.29\text{e-}03$	0.084 ± 0.0030	1.95 ± 1.07
		$-2.10\text{e-}03 \pm 2.10\text{e-}03$	$5.89\text{e-}03 \pm 1.97\text{e-}03$	0.073 ± 0.0043	16.79 ± 1.80
		$-4.67\text{e-}03 \pm 1.67\text{e-}02$	$8.08\text{e-}03 \pm 1.62\text{e-}02$	0.083 ± 0.0315	0.42 ± 11.06
	
LFC 2180	0.55
		$1.52\text{e-}02 \pm 3.04\text{e-}03$	$3.47\text{e-}03 \pm 2.05\text{e-}03$	0.128 ± 0.0037	36.23 ± 0.88
		$-4.77\text{e-}03 \pm 1.56\text{e-}02$	$2.49\text{e-}02 \pm 1.51\text{e-}02$	0.190 ± 0.0287	30.93 ± 4.76
	
LFC 2181	0.60	$-1.63\text{e-}03 \pm 1.11\text{e-}03$	$-1.76\text{e-}03 \pm 1.06\text{e-}03$	0.058 ± 0.0023	1.93 ± 1.21
		$3.11\text{e-}03 \pm 1.10\text{e-}03$	$-3.54\text{e-}03 \pm 1.05\text{e-}03$	0.046 ± 0.0023	35.07 ± 1.50
		$-1.20\text{e-}02 \pm 1.31\text{e-}02$	$-3.89\text{e-}02 \pm 1.43\text{e-}02$	0.036 ± 0.0294	2.69 ± 22.67
	
LFC 2184	0.54	$1.24\text{e-}02 \pm 4.70\text{e-}04$	$2.80\text{e-}03 \pm 3.47\text{e-}04$	0.161 ± 0.0014	4.26 ± 0.29
		$6.59\text{e-}03 \pm 1.12\text{e-}03$	$2.34\text{e-}03 \pm 1.01\text{e-}03$	0.113 ± 0.0023	3.42 ± 0.61
		$-1.49\text{e-}02 \pm 1.94\text{e-}03$	$1.15\text{e-}03 \pm 1.70\text{e-}03$	0.205 ± 0.0038	4.79 ± 0.50
		$-1.66\text{e-}02 \pm 5.87\text{e-}03$	$-2.84\text{e-}04 \pm 5.20\text{e-}03$	0.305 ± 0.0119	0.00 ± 0.91
LFC 2310	0.61	$3.67\text{e-}03 \pm 1.21\text{e-}03$	$8.02\text{e-}03 \pm 9.49\text{e-}04$	0.307 ± 0.0022	0.64 ± 0.26
		$5.30\text{e-}03 \pm 1.37\text{e-}03$	$7.34\text{e-}03 \pm 1.26\text{e-}03$	0.315 ± 0.0025	0.77 ± 0.27
	
	
LFC 2356	0.73	$-1.70\text{e-}02 \pm 2.87\text{e-}03$	$1.89\text{e-}02 \pm 2.62\text{e-}03$	0.397 ± 0.0047	5.72 ± 0.43
		$-4.19\text{e-}03 \pm 4.79\text{e-}03$	$-6.09\text{e-}04 \pm 4.02\text{e-}03$	0.430 ± 0.0048	1.20 ± 0.42
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a^3}{a} \rangle$ (3)	$\langle \frac{a^4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 2362	0.65	$1.30\text{e-}03 \pm 2.76\text{e-}03$	$3.51\text{e-}03 \pm 2.65\text{e-}03$	0.104 ± 0.0055	7.36 ± 1.57
		$-3.98\text{e-}03 \pm 3.17\text{e-}03$	$-1.14\text{e-}02 \pm 3.03\text{e-}03$	0.141 ± 0.0061	14.61 ± 1.33
		$-2.34\text{e-}02 \pm 9.39\text{e-}03$	$3.40\text{e-}02 \pm 8.84\text{e-}03$	0.189 ± 0.0160	1.34 ± 2.33
		$-5.85\text{e-}02 \pm 2.52\text{e-}02$	$5.92\text{e-}02 \pm 2.40\text{e-}02$	0.377 ± 0.0378	0.00 ± 3.13
LFC 2405	0.51	$-2.20\text{e-}02 \pm 4.63\text{e-}03$	$-2.76\text{e-}03 \pm 4.04\text{e-}03$	0.117 ± 0.0095	37.59 ± 2.50
		$-3.56\text{e-}02 \pm 3.38\text{e-}02$	$1.48\text{e-}02 \pm 3.15\text{e-}02$	0.170 ± 0.0649	0.00 ± 10.86
	
	
LFC 2414	0.64	$2.63\text{e-}03 \pm 5.81\text{e-}04$	$5.43\text{e-}03 \pm 4.76\text{e-}04$	0.056 ± 0.0012	12.52 ± 0.63
		$4.18\text{e-}03 \pm 1.22\text{e-}03$	$-1.24\text{e-}03 \pm 1.19\text{e-}03$	0.070 ± 0.0025	9.31 ± 1.05
		$1.78\text{e-}03 \pm 4.34\text{e-}03$	$1.50\text{e-}02 \pm 4.34\text{e-}03$	0.114 ± 0.0108	7.53 ± 2.41
	
LFC 2422	0.96	$1.39\text{e-}02 \pm 8.44\text{e-}04$	$1.78\text{e-}03 \pm 7.42\text{e-}04$	0.107 ± 0.0022	5.13 ± 0.60
		$1.04\text{e-}02 \pm 1.23\text{e-}03$	$1.92\text{e-}03 \pm 1.15\text{e-}03$	0.142 ± 0.0026	13.06 ± 0.57
		$-3.48\text{e-}02 \pm 1.06\text{e-}02$	$1.22\text{e-}02 \pm 9.93\text{e-}03$	0.221 ± 0.0180	3.15 ± 2.61
	
LFC 2428	0.68	$-4.07\text{e-}03 \pm 2.71\text{e-}03$	$-3.75\text{e-}03 \pm 2.64\text{e-}03$	0.058 ± 0.0060	0.05 ± 3.09
		$-7.75\text{e-}04 \pm 1.86\text{e-}03$	$1.47\text{e-}03 \pm 1.80\text{e-}03$	0.069 ± 0.0039	19.57 ± 1.54
		$-9.31\text{e-}03 \pm 8.30\text{e-}03$	$-4.02\text{e-}03 \pm 8.02\text{e-}03$	0.215 ± 0.0183	11.05 ± 2.33
	
LFC 2472	0.89	$1.97\text{e-}03 \pm 1.88\text{e-}03$	$-1.91\text{e-}03 \pm 1.82\text{e-}03$	0.223 ± 0.0037	0.69 ± 0.53
		$-2.67\text{e-}02 \pm 5.15\text{e-}03$	$1.74\text{e-}02 \pm 4.65\text{e-}03$	0.260 ± 0.0093	15.52 ± 1.11
	
	
LFC 2484	0.94	$4.25\text{e-}03 \pm 2.68\text{e-}03$	$-1.34\text{e-}03 \pm 2.53\text{e-}03$	0.013 ± 0.0068	47.87 ± 14.27
		$-1.10\text{e-}02 \pm 3.78\text{e-}03$	$-1.36\text{e-}03 \pm 3.53\text{e-}03$	0.029 ± 0.0078	6.98 ± 5.43
		$-4.92\text{e-}02 \pm 3.67\text{e-}02$	$3.83\text{e-}02 \pm 3.46\text{e-}02$	0.109 ± 0.0657	0.00 ± 18.27
	
LFC 2503	0.88
		$-1.49\text{e-}02 \pm 1.99\text{e-}03$	$-1.50\text{e-}03 \pm 1.92\text{e-}03$	0.129 ± 0.0043	2.85 ± 1.02
		$1.46\text{e-}02 \pm 7.49\text{e-}03$	$4.16\text{e-}03 \pm 7.85\text{e-}03$	0.088 ± 0.0155	0.02 ± 4.86
	
LFC 2525	0.88	$-1.66\text{e-}02 \pm 1.23\text{e-}03$	$6.48\text{e-}03 \pm 1.01\text{e-}03$	0.347 ± 0.0027	0.46 ± 0.28
		$-1.90\text{e-}02 \pm 3.45\text{e-}03$	$1.47\text{e-}02 \pm 3.19\text{e-}03$	0.279 ± 0.0062	2.55 ± 0.78
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a^3}{a} \rangle$ (3)	$\langle \frac{a^4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 2539	0.62	$-5.41\text{e-}03 \pm 7.18\text{e-}03$	$-1.42\text{e-}02 \pm 6.28\text{e-}03$	0.106 ± 0.0147	0.00 ± 4.32
		$-4.39\text{e-}03 \pm 2.44\text{e-}03$	$9.42\text{e-}04 \pm 2.28\text{e-}03$	0.098 ± 0.0050	10.37 ± 1.55
		$7.38\text{e-}02 \pm 1.51\text{e-}02$	$-2.73\text{e-}02 \pm 1.13\text{e-}02$	0.151 ± 0.0238	41.12 ± 4.51
		$9.06\text{e-}02 \pm 4.79\text{e-}02$	$-1.57\text{e-}02 \pm 2.44\text{e-}02$	0.264 ± 0.0509	9.70 ± 5.54
LFC 2549	0.51	$8.29\text{e-}05 \pm 4.58\text{e-}04$	$3.88\text{e-}03 \pm 3.52\text{e-}04$	0.109 ± 0.0011	0.01 ± 0.30
		$-4.03\text{e-}03 \pm 4.80\text{e-}04$	$5.78\text{e-}04 \pm 4.55\text{e-}04$	0.158 ± 0.0010	1.78 ± 0.19
		$6.66\text{e-}04 \pm 2.13\text{e-}03$	$2.10\text{e-}02 \pm 1.95\text{e-}03$	0.265 ± 0.0038	7.37 ± 0.47
		$-1.81\text{e-}02 \pm 8.26\text{e-}03$	$1.27\text{e-}02 \pm 7.78\text{e-}03$	0.177 ± 0.0143	13.43 ± 2.58
LFC 2568	0.62	$-1.99\text{e-}03 \pm 2.71\text{e-}03$	$-1.52\text{e-}03 \pm 2.62\text{e-}03$	0.032 ± 0.0058	14.71 ± 5.22
		$2.68\text{e-}02 \pm 4.57\text{e-}03$	$-4.00\text{e-}03 \pm 3.63\text{e-}03$	0.103 ± 0.0093	32.28 ± 1.83
		$9.02\text{e-}02 \pm 2.30\text{e-}02$	$-4.00\text{e-}03 \pm 1.38\text{e-}02$	0.399 ± 0.0239	5.12 ± 2.18
	
LFC 2670	0.62	$9.64\text{e-}03 \pm 2.33\text{e-}03$	$5.95\text{e-}03 \pm 2.22\text{e-}03$	0.141 ± 0.0063	3.24 ± 1.41
		$-6.90\text{e-}03 \pm 3.42\text{e-}03$	$4.56\text{e-}03 \pm 2.55\text{e-}03$	0.068 ± 0.0080	41.26 ± 3.72
	
	
LFC 2692	0.57	$1.57\text{e-}03 \pm 3.15\text{e-}03$	$6.35\text{e-}03 \pm 2.95\text{e-}03$	0.183 ± 0.0064	13.41 ± 1.12
		$2.95\text{e-}02 \pm 5.26\text{e-}02$	$-6.54\text{e-}03 \pm 4.76\text{e-}02$	0.250 ± 0.0836	0.00 ± 15.23
	
	
LFC 2708	0.68	$-1.14\text{e-}02 \pm 2.15\text{e-}03$	$-3.11\text{e-}03 \pm 2.06\text{e-}03$	0.027 ± 0.0081	2.91 ± 8.66
		$-8.51\text{e-}03 \pm 1.92\text{e-}03$	$2.44\text{e-}03 \pm 1.82\text{e-}03$	0.040 ± 0.0049	4.34 ± 2.99
		$-4.14\text{e-}02 \pm 1.13\text{e-}02$	$5.05\text{e-}03 \pm 6.96\text{e-}03$	0.180 ± 0.0121	2.50 ± 2.14
	
LFC 2709	0.71	$-1.06\text{e-}02 \pm 2.79\text{e-}03$	$1.43\text{e-}02 \pm 2.34\text{e-}03$	0.192 ± 0.0056	3.21 ± 0.94
		$-1.38\text{e-}02 \pm 4.14\text{e-}03$	$-1.73\text{e-}02 \pm 4.05\text{e-}03$	0.239 ± 0.0074	4.97 ± 0.97
	
	
LFC 2712	0.55	$-4.37\text{e-}05 \pm 8.38\text{e-}04$	$9.33\text{e-}03 \pm 6.08\text{e-}04$	0.343 ± 0.0016	2.79 ± 0.17
		$-1.39\text{e-}02 \pm 1.41\text{e-}03$	$1.31\text{e-}02 \pm 1.15\text{e-}03$	0.355 ± 0.0027	1.11 ± 0.29
		$3.53\text{e-}03 \pm 4.21\text{e-}03$	$1.49\text{e-}02 \pm 3.99\text{e-}03$	0.279 ± 0.0091	11.27 ± 1.04
	
LFC 2719	0.67	$3.84\text{e-}03 \pm 2.28\text{e-}03$	$-4.21\text{e-}03 \pm 2.05\text{e-}03$	0.046 ± 0.0049	30.63 ± 3.15
		$7.28\text{e-}03 \pm 3.44\text{e-}03$	$-4.65\text{e-}03 \pm 3.52\text{e-}03$	0.193 ± 0.0078	29.83 ± 1.35
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 2742	0.83	$1.67\text{e-}03 \pm 6.35\text{e-}04$	$3.32\text{e-}03 \pm 4.22\text{e-}04$	0.101 ± 0.0013	5.68 ± 0.37
		$-2.26\text{e-}03 \pm 6.20\text{e-}04$	$8.44\text{e-}03 \pm 4.16\text{e-}04$	0.159 ± 0.0013	4.47 ± 0.22
		$-3.60\text{e-}03 \pm 1.11\text{e-}03$	$-2.46\text{e-}03 \pm 1.09\text{e-}03$	0.258 ± 0.0023	3.11 ± 0.26
		$-2.88\text{e-}03 \pm 2.82\text{e-}03$	$-4.38\text{e-}03 \pm 2.76\text{e-}03$	0.352 ± 0.0054	0.26 ± 0.53
LFC 2782	0.64	$1.04\text{e-}03 \pm 7.70\text{e-}04$	$1.06\text{e-}02 \pm 5.19\text{e-}04$	0.445 ± 0.0013	0.55 ± 0.11
		$2.23\text{e-}03 \pm 6.04\text{e-}04$	$9.36\text{e-}03 \pm 4.77\text{e-}04$	0.512 ± 0.0010	1.18 ± 0.08
		$3.35\text{e-}03 \pm 1.17\text{e-}03$	$4.46\text{e-}03 \pm 1.13\text{e-}03$	0.504 ± 0.0017	0.32 ± 0.14
		$5.56\text{e-}03 \pm 5.64\text{e-}03$	$1.08\text{e-}02 \pm 5.86\text{e-}03$	0.486 ± 0.0086	1.94 ± 0.66
LFC 2888	0.77
		$-5.39\text{e-}03 \pm 1.26\text{e-}03$	$-8.62\text{e-}03 \pm 1.11\text{e-}03$	0.068 ± 0.0031	5.39 ± 1.38
		$2.22\text{e-}02 \pm 5.12\text{e-}03$	$-8.92\text{e-}03 \pm 4.93\text{e-}03$	0.107 ± 0.0101	16.88 ± 2.81
		$5.86\text{e-}04 \pm 8.92\text{e-}03$	$1.47\text{e-}02 \pm 1.46\text{e-}02$	0.147 ± 0.0169	0.00 ± 3.57
LFC 2955	0.48
		$3.76\text{e-}03 \pm 1.16\text{e-}03$	$-4.20\text{e-}03 \pm 1.06\text{e-}03$	0.213 ± 0.0025	0.98 ± 0.38
		$2.72\text{e-}03 \pm 4.54\text{e-}03$	$3.73\text{e-}03 \pm 4.39\text{e-}03$	0.179 ± 0.0087	7.41 ± 1.52
		$1.32\text{e-}02 \pm 1.20\text{e-}02$	$2.78\text{e-}03 \pm 6.18\text{e-}03$	0.199 ± 0.0116	0.00 ± 1.87
LFC 2956	0.98	$-6.19\text{e-}03 \pm 5.98\text{e-}03$	$-6.54\text{e-}03 \pm 5.27\text{e-}03$	0.181 ± 0.0116	0.00 ± 2.03
		$-7.68\text{e-}03 \pm 2.12\text{e-}03$	$-1.36\text{e-}02 \pm 1.92\text{e-}03$	0.168 ± 0.0041	9.16 ± 0.79
		$-1.54\text{e-}03 \pm 5.12\text{e-}03$	$-2.60\text{e-}02 \pm 7.53\text{e-}03$	0.157 ± 0.0096	0.60 ± 1.87
	
LFC 2988	0.82	$3.66\text{e-}03 \pm 3.02\text{e-}03$	$-1.00\text{e-}02 \pm 2.64\text{e-}03$	0.081 ± 0.0080	0.78 ± 2.99
		$-5.16\text{e-}03 \pm 1.97\text{e-}03$	$-7.03\text{e-}03 \pm 1.44\text{e-}03$	0.073 ± 0.0054	21.36 ± 2.24
		$-4.94\text{e-}03 \pm 8.08\text{e-}03$	$7.18\text{e-}03 \pm 7.96\text{e-}03$	0.084 ± 0.0154	159.80 ± 5.44
		$2.50\text{e-}02 \pm 1.12\text{e-}02$	$2.10\text{e-}02 \pm 1.09\text{e-}02$	0.230 ± 0.0208	149.30 ± 2.88
LFC 3026	0.72
		$2.64\text{e-}03 \pm 1.28\text{e-}03$	$-5.32\text{e-}03 \pm 1.03\text{e-}03$	0.127 ± 0.0027	44.32 ± 0.69
		$-2.08\text{e-}02 \pm 3.83\text{e-}03$	$1.58\text{e-}02 \pm 3.50\text{e-}03$	0.143 ± 0.0079	1.26 ± 1.64
		$-1.69\text{e-}02 \pm 7.99\text{e-}03$	$7.60\text{e-}03 \pm 7.77\text{e-}03$	0.218 ± 0.0141	38.36 ± 2.09
LFC 3072	0.77
		$1.20\text{e-}03 \pm 2.29\text{e-}03$	$-5.28\text{e-}03 \pm 2.24\text{e-}03$	0.069 ± 0.0047	7.32 ± 2.12
		$-1.62\text{e-}02 \pm 8.89\text{e-}03$	$1.70\text{e-}03 \pm 9.03\text{e-}03$	0.132 ± 0.0165	5.11 ± 3.22
		$-7.83\text{e-}02 \pm 2.78\text{e-}02$	$3.80\text{e-}02 \pm 2.17\text{e-}02$	0.242 ± 0.0383	12.70 ± 5.15
LFC 3087	0.68
		$4.43\text{e-}03 \pm 1.13\text{e-}03$	$-1.23\text{e-}02 \pm 7.13\text{e-}04$	0.095 ± 0.0024	0.68 ± 0.77
		$1.60\text{e-}02 \pm 3.37\text{e-}03$	$-5.19\text{e-}03 \pm 3.15\text{e-}03$	0.075 ± 0.0070	27.24 ± 2.71
		$-1.64\text{e-}03 \pm 6.69\text{e-}03$	$-7.12\text{e-}03 \pm 6.83\text{e-}03$	0.058 ± 0.0132	17.35 ± 6.13

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a^3}{a} \rangle$ (3)	$\langle \frac{a^4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 3094	0.67	$2.38\text{e-}03 \pm 1.04\text{e-}03$	$1.86\text{e-}03 \pm 1.01\text{e-}03$	0.088 ± 0.0024	0.16 ± 0.81
		$5.99\text{e-}03 \pm 7.34\text{e-}04$	$-5.83\text{e-}04 \pm 6.93\text{e-}04$	0.082 ± 0.0016	4.38 ± 0.61
		$-1.23\text{e-}02 \pm 3.50\text{e-}03$	$9.65\text{e-}03 \pm 3.31\text{e-}03$	0.101 ± 0.0074	10.53 ± 2.32
		$1.56\text{e-}03 \pm 1.30\text{e-}02$	$3.49\text{e-}04 \pm 1.30\text{e-}02$	0.125 ± 0.0257	14.59 ± 5.50
LFC 3218	0.59
		$1.66\text{e-}03 \pm 1.45\text{e-}03$	$7.00\text{e-}03 \pm 1.31\text{e-}03$	0.120 ± 0.0031	4.06 ± 0.82
		$-1.65\text{e-}02 \pm 4.11\text{e-}03$	$-1.64\text{e-}03 \pm 3.85\text{e-}03$	0.085 ± 0.0086	3.89 ± 3.09
		$-4.45\text{e-}03 \pm 7.09\text{e-}03$	$1.81\text{e-}02 \pm 7.23\text{e-}03$	0.210 ± 0.0134	27.80 ± 2.08
LFC 3274	0.75	$-2.15\text{e-}02 \pm 3.81\text{e-}03$	$-1.81\text{e-}02 \pm 3.25\text{e-}03$	0.218 ± 0.0079	1.10 ± 1.19
		$-1.92\text{e-}03 \pm 7.80\text{e-}04$	$-2.09\text{e-}04 \pm 5.05\text{e-}04$	0.185 ± 0.0013	78.36 ± 0.34
	
	
LFC 3292	0.59	$1.18\text{e-}02 \pm 2.08\text{e-}03$	$5.03\text{e-}03 \pm 1.95\text{e-}03$	0.169 ± 0.0049	1.11 ± 0.92
		$9.95\text{e-}04 \pm 1.37\text{e-}03$	$3.75\text{e-}03 \pm 1.30\text{e-}03$	0.203 ± 0.0027	4.92 ± 0.40
		$-3.29\text{e-}02 \pm 4.87\text{e-}03$	$-2.69\text{e-}02 \pm 4.48\text{e-}03$	0.312 ± 0.0088	0.43 ± 0.84
		$-6.19\text{e-}03 \pm 8.97\text{e-}03$	$-1.71\text{e-}02 \pm 8.89\text{e-}03$	0.334 ± 0.0180	1.24 ± 1.59
LFC 3329	0.54
		$-5.89\text{e-}03 \pm 1.49\text{e-}03$	$1.34\text{e-}02 \pm 1.12\text{e-}03$	0.208 ± 0.0030	4.30 ± 0.46
		$-4.96\text{e-}03 \pm 4.06\text{e-}03$	$3.33\text{e-}02 \pm 3.83\text{e-}03$	0.320 ± 0.0069	0.95 ± 0.75
		$1.15\text{e-}02 \pm 6.84\text{e-}03$	$4.12\text{e-}02 \pm 6.92\text{e-}03$	0.368 ± 0.0105	4.72 ± 1.01
LFC 3359	0.51
		$4.39\text{e-}04 \pm 1.50\text{e-}03$	$-7.17\text{e-}03 \pm 1.27\text{e-}03$	0.110 ± 0.0033	8.39 ± 0.92
		$-2.26\text{e-}03 \pm 5.87\text{e-}03$	$5.83\text{e-}03 \pm 5.78\text{e-}03$	0.157 ± 0.0110	13.07 ± 2.15
		$-4.00\text{e-}02 \pm 5.30\text{e-}02$	$-2.89\text{e-}02 \pm 4.79\text{e-}02$	0.175 ± 0.0787	0.00 ± 14.11
LFC 3381	0.66
		$1.21\text{e-}02 \pm 1.48\text{e-}03$	$1.07\text{e-}03 \pm 1.36\text{e-}03$	0.093 ± 0.0037	10.84 ± 1.20
		$9.62\text{e-}03 \pm 5.81\text{e-}03$	$-1.25\text{e-}03 \pm 5.44\text{e-}03$	0.072 ± 0.0118	26.44 ± 4.57
		$3.24\text{e-}02 \pm 1.74\text{e-}02$	$1.27\text{e-}02 \pm 1.62\text{e-}02$	0.149 ± 0.0295	98.18 ± 6.13
LFC 3395	0.63
		$-1.16\text{e-}03 \pm 8.63\text{e-}04$	$3.41\text{e-}03 \pm 7.85\text{e-}04$	0.147 ± 0.0018	1.75 ± 0.38
		$5.18\text{e-}03 \pm 2.45\text{e-}03$	$6.13\text{e-}03 \pm 2.37\text{e-}03$	0.151 ± 0.0047	2.10 ± 0.97
		$2.52\text{e-}02 \pm 6.33\text{e-}03$	$1.54\text{e-}02 \pm 5.93\text{e-}03$	0.152 ± 0.0123	27.61 ± 2.35
LFC 3423	0.67
		$-3.85\text{e-}03 \pm 1.15\text{e-}03$	$6.24\text{e-}03 \pm 1.27\text{e-}03$	0.170 ± 0.0018	5.29 ± 0.34
		$4.31\text{e-}02 \pm 1.05\text{e-}02$	$1.81\text{e-}02 \pm 8.62\text{e-}03$	0.277 ± 0.0174	11.56 ± 2.13
		$6.00\text{e-}02 \pm 6.08\text{e-}02$	$8.30\text{e-}03 \pm 4.81\text{e-}02$	0.363 ± 0.0794	0.00 ± 7.85

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a^3}{a} \rangle$ (3)	$\langle \frac{a^4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 3427	0.47	$-3.12\text{e-}02 \pm 8.43\text{e-}03$	$-3.91\text{e-}02 \pm 7.01\text{e-}03$	0.320 ± 0.0179	3.07 ± 1.98
		$-4.24\text{e-}02 \pm 3.94\text{e-}03$	$-2.85\text{e-}02 \pm 2.86\text{e-}03$	0.251 ± 0.0077	19.69 ± 1.05
		$-2.99\text{e-}03 \pm 6.46\text{e-}03$	$7.05\text{e-}03 \pm 8.17\text{e-}03$	0.232 ± 0.0108	0.00 ± 1.54
	
LFC 3430	0.51	$2.02\text{e-}03 \pm 3.08\text{e-}03$	$2.30\text{e-}02 \pm 1.60\text{e-}03$	0.286 ± 0.0055	0.25 ± 0.66
		$-2.96\text{e-}03 \pm 1.92\text{e-}03$	$3.23\text{e-}02 \pm 1.13\text{e-}03$	0.365 ± 0.0032	1.50 ± 0.31
		$-1.35\text{e-}03 \pm 3.40\text{e-}03$	$5.31\text{e-}02 \pm 3.19\text{e-}03$	0.515 ± 0.0048	0.28 ± 0.38
		$8.02\text{e-}03 \pm 3.26\text{e-}03$	$5.02\text{e-}02 \pm 3.15\text{e-}03$	0.528 ± 0.0045	5.08 ± 0.35
LFC 3458	0.63
		$-5.68\text{e-}03 \pm 9.79\text{e-}04$	$1.64\text{e-}04 \pm 9.02\text{e-}04$	0.063 ± 0.0023	10.08 ± 1.10
		$-2.03\text{e-}02 \pm 4.08\text{e-}03$	$4.92\text{e-}03 \pm 3.85\text{e-}03$	0.026 ± 0.0086	23.91 ± 9.08
		$-5.25\text{e-}02 \pm 1.81\text{e-}02$	$4.74\text{e-}02 \pm 1.62\text{e-}02$	0.072 ± 0.0310	1.97 ± 12.31
LFC 3508	0.78	$-9.73\text{e-}04 \pm 9.53\text{e-}04$	$3.77\text{e-}03 \pm 7.77\text{e-}04$	0.121 ± 0.0020	0.99 ± 0.50
		$1.05\text{e-}03 \pm 9.55\text{e-}04$	$7.10\text{e-}03 \pm 6.59\text{e-}04$	0.126 ± 0.0019	84.05 ± 0.46
		$8.77\text{e-}02 \pm 6.95\text{e-}03$	$5.97\text{e-}02 \pm 5.08\text{e-}03$	0.152 ± 0.0119	2.91 ± 2.50
		$3.69\text{e-}02 \pm 7.11\text{e-}03$	$3.18\text{e-}02 \pm 6.52\text{e-}03$	0.108 ± 0.0138	1.68 ± 3.67
LFC 3562	0.51
		$-1.91\text{e-}03 \pm 1.50\text{e-}03$	$8.12\text{e-}04 \pm 1.34\text{e-}03$	0.039 ± 0.0034	1.47 ± 2.38
		$1.35\text{e-}03 \pm 4.10\text{e-}03$	$3.62\text{e-}03 \pm 3.58\text{e-}03$	0.060 ± 0.0084	9.35 ± 4.11
		$-2.72\text{e-}02 \pm 1.00\text{e-}02$	$1.28\text{e-}02 \pm 1.01\text{e-}02$	0.049 ± 0.0186	7.71 ± 8.79
LFC 3577	0.76	$-5.12\text{e-}02 \pm 5.09\text{e-}03$	$1.88\text{e-}03 \pm 1.46\text{e-}03$	0.072 ± 0.0160	0.11 ± 6.71
		$6.43\text{e-}03 \pm 1.96\text{e-}03$	$4.56\text{e-}03 \pm 1.08\text{e-}03$	0.033 ± 0.0063	27.54 ± 4.21
		$7.87\text{e-}03 \pm 6.81\text{e-}03$	$-1.57\text{e-}03 \pm 6.63\text{e-}03$	0.088 ± 0.0140	40.58 ± 4.60
		$-3.44\text{e-}02 \pm 1.58\text{e-}02$	$-1.11\text{e-}03 \pm 1.29\text{e-}02$	0.140 ± 0.0252	5.96 ± 5.38
LFC 3632	0.59
		$-1.09\text{e-}02 \pm 1.67\text{e-}03$	$-4.28\text{e-}03 \pm 1.56\text{e-}03$	0.126 ± 0.0039	3.50 ± 0.95
		$-3.63\text{e-}04 \pm 6.96\text{e-}04$	$-1.31\text{e-}05 \pm 6.46\text{e-}04$	0.038 ± 0.0018	55.74 ± 1.94
	
LFC 3646	0.50
		$-9.53\text{e-}03 \pm 1.59\text{e-}03$	$-4.94\text{e-}03 \pm 1.36\text{e-}03$	0.096 ± 0.0040	18.55 ± 1.23
		$9.27\text{e-}03 \pm 4.05\text{e-}03$	$1.85\text{e-}02 \pm 3.84\text{e-}03$	0.218 ± 0.0052	4.40 ± 0.72
	
LFC 3760	0.91	$1.29\text{e-}02 \pm 2.22\text{e-}03$	$-2.50\text{e-}03 \pm 1.70\text{e-}03$	0.084 ± 0.0084	9.95 ± 2.98
		$-1.98\text{e-}02 \pm 2.61\text{e-}03$	$-1.74\text{e-}02 \pm 1.97\text{e-}03$	0.055 ± 0.0061	34.67 ± 3.48
	
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 3808	0.49
		$7.50\text{e-}03 \pm 1.28\text{e-}03$	$-1.81\text{e-}03 \pm 9.83\text{e-}04$	0.117 ± 0.0028	2.44 ± 0.76
		$-1.37\text{e-}02 \pm 6.64\text{e-}03$	$-2.59\text{e-}03 \pm 6.42\text{e-}03$	0.109 ± 0.0134	4.91 ± 3.41
		$4.45\text{e-}02 \pm 2.90\text{e-}02$	$9.69\text{e-}03 \pm 2.46\text{e-}02$	0.235 ± 0.0479	2.10 ± 6.21
LFC 3878	0.65
		$8.43\text{e-}03 \pm 1.79\text{e-}03$	$9.09\text{e-}05 \pm 1.71\text{e-}03$	0.187 ± 0.0037	6.38 ± 0.64
		$-8.32\text{e-}03 \pm 1.29\text{e-}02$	$3.26\text{e-}02 \pm 9.77\text{e-}03$	0.234 ± 0.0134	19.88 ± 1.84
	
LFC 3894	0.82
		$6.27\text{e-}03 \pm 1.38\text{e-}03$	$-5.61\text{e-}05 \pm 1.32\text{e-}03$	0.159 ± 0.0028	0.40 ± 0.56
		$-3.03\text{e-}03 \pm 5.14\text{e-}03$	$4.70\text{e-}03 \pm 4.55\text{e-}03$	0.112 ± 0.0078	3.64 ± 2.03
	
LFC 3912	0.79
		$2.87\text{e-}02 \pm 2.02\text{e-}03$	$-1.08\text{e-}02 \pm 1.44\text{e-}03$	0.135 ± 0.0054	0.62 ± 1.22
		$2.50\text{e-}02 \pm 6.88\text{e-}03$	$1.76\text{e-}03 \pm 6.28\text{e-}03$	0.105 ± 0.0142	3.03 ± 4.26
		$-8.06\text{e-}03 \pm 1.76\text{e-}02$	$8.94\text{e-}03 \pm 1.76\text{e-}02$	0.123 ± 0.0328	19.31 ± 8.48
LFC 3921	0.86	$3.22\text{e-}03 \pm 2.73\text{e-}03$	$-4.54\text{e-}03 \pm 2.58\text{e-}03$	0.063 ± 0.0060	38.79 ± 2.83
		$8.81\text{e-}03 \pm 7.77\text{e-}03$	$-1.08\text{e-}03 \pm 7.43\text{e-}03$	0.124 ± 0.0152	125.20 ± 3.16
	
	
LFC 3927	0.50
		$1.19\text{e-}02 \pm 7.45\text{e-}04$	$1.03\text{e-}03 \pm 7.04\text{e-}04$	0.219 ± 0.0019	7.48 ± 0.29
		$1.83\text{e-}02 \pm 4.14\text{e-}03$	$-1.23\text{e-}02 \pm 4.01\text{e-}03$	0.171 ± 0.0080	15.68 ± 1.48
		$-1.97\text{e-}02 \pm 1.12\text{e-}02$	$3.42\text{e-}02 \pm 1.13\text{e-}02$	0.185 ± 0.0199	3.60 ± 3.09
LFC 3942	0.71
		$-1.67\text{e-}02 \pm 9.42\text{e-}04$	$-8.94\text{e-}03 \pm 5.97\text{e-}04$	0.267 ± 0.0025	7.09 ± 0.32
		$2.89\text{e-}02 \pm 1.46\text{e-}03$	$-8.96\text{e-}03 \pm 8.20\text{e-}04$	0.229 ± 0.0046	2.73 ± 0.68
		$4.35\text{e-}02 \pm 3.32\text{e-}03$	$-6.64\text{e-}03 \pm 2.56\text{e-}03$	0.113 ± 0.0084	49.38 ± 2.34
LFC 4009	0.61
		$-2.32\text{e-}02 \pm 1.08\text{e-}03$	$5.01\text{e-}03 \pm 8.89\text{e-}04$	0.196 ± 0.0035	0.67 ± 0.57
		$2.59\text{e-}03 \pm 3.40\text{e-}03$	$-3.41\text{e-}03 \pm 3.31\text{e-}03$	0.183 ± 0.0069	11.85 ± 1.22
		$-2.60\text{e-}02 \pm 1.29\text{e-}02$	$-2.73\text{e-}03 \pm 1.22\text{e-}02$	0.167 ± 0.0228	6.60 ± 4.04
LFC 4029	0.89	$-3.97\text{e-}03 \pm 4.39\text{e-}03$	$1.40\text{e-}02 \pm 3.77\text{e-}03$	0.318 ± 0.0085	0.11 ± 0.94
		$-2.54\text{e-}03 \pm 1.40\text{e-}03$	$7.26\text{e-}03 \pm 1.20\text{e-}03$	0.350 ± 0.0025	2.79 ± 0.25
		$3.23\text{e-}03 \pm 4.57\text{e-}03$	$-9.97\text{e-}03 \pm 4.47\text{e-}03$	0.337 ± 0.0077	18.98 ± 0.81
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a3}{a} \rangle$ (3)	$\langle \frac{a4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 4059	0.68	$7.78\text{e-}03 \pm 1.07\text{e-}03$	$2.58\text{e-}05 \pm 1.03\text{e-}03$	0.248 ± 0.0026	3.52 ± 0.35
		$8.21\text{e-}03 \pm 1.37\text{e-}03$	$-5.22\text{e-}03 \pm 1.28\text{e-}03$	0.254 ± 0.0033	7.74 ± 0.44
		$-5.13\text{e-}02 \pm 7.30\text{e-}03$	$-1.43\text{e-}02 \pm 6.23\text{e-}03$	0.214 ± 0.0124	35.81 ± 2.17
		$2.22\text{e-}02 \pm 1.25\text{e-}02$	$-1.82\text{e-}02 \pm 1.20\text{e-}02$	0.180 ± 0.0206	45.55 ± 3.64
LFC 4095	0.63
		$1.10\text{e-}02 \pm 1.62\text{e-}03$	$2.08\text{e-}03 \pm 1.51\text{e-}03$	0.176 ± 0.0033	5.12 ± 0.59
		$1.14\text{e-}02 \pm 4.21\text{e-}03$	$1.63\text{e-}02 \pm 4.03\text{e-}03$	0.206 ± 0.0081	7.65 ± 1.27
		$1.55\text{e-}02 \pm 1.64\text{e-}02$	$1.30\text{e-}02 \pm 1.57\text{e-}02$	0.177 ± 0.0316	14.40 ± 6.61
LFC 4188	0.66
		$-7.68\text{e-}03 \pm 1.04\text{e-}03$	$1.23\text{e-}02 \pm 7.89\text{e-}04$	0.265 ± 0.0021	1.72 ± 0.26
		$-5.01\text{e-}04 \pm 2.75\text{e-}03$	$1.12\text{e-}02 \pm 2.51\text{e-}03$	0.207 ± 0.0050	4.55 ± 0.81
		$1.34\text{e-}02 \pm 1.09\text{e-}02$	$5.50\text{e-}03 \pm 1.05\text{e-}02$	0.095 ± 0.0210	10.10 ± 6.65
LFC 4211	0.63
		$-1.12\text{e-}02 \pm 1.15\text{e-}03$	$1.36\text{e-}02 \pm 7.54\text{e-}04$	0.225 ± 0.0024	1.38 ± 0.34
		$-1.41\text{e-}02 \pm 1.64\text{e-}03$	$5.30\text{e-}03 \pm 1.52\text{e-}03$	0.247 ± 0.0033	1.23 ± 0.45
		$-1.46\text{e-}02 \pm 5.45\text{e-}03$	$-1.07\text{e-}02 \pm 5.26\text{e-}03$	0.237 ± 0.0097	0.48 ± 1.34
LFC 4248	0.49	$9.88\text{e-}04 \pm 1.37\text{e-}03$	$9.61\text{e-}03 \pm 1.11\text{e-}03$	0.530 ± 0.0021	1.76 ± 0.17
		$-5.65\text{e-}03 \pm 1.35\text{e-}03$	$1.85\text{e-}02 \pm 1.07\text{e-}03$	0.575 ± 0.0019	0.53 ± 0.14
		$1.26\text{e-}02 \pm 2.19\text{e-}03$	$1.11\text{e-}02 \pm 2.00\text{e-}03$	0.589 ± 0.0030	0.66 ± 0.23
		$-8.03\text{e-}03 \pm 2.04\text{e-}02$	$-2.81\text{e-}03 \pm 2.16\text{e-}02$	0.499 ± 0.0290	7.42 ± 2.41
LFC 4394	0.84
		$-1.56\text{e-}02 \pm 1.96\text{e-}03$	$4.11\text{e-}03 \pm 1.84\text{e-}03$	0.197 ± 0.0044	10.71 ± 0.72
		$2.76\text{e-}03 \pm 5.02\text{e-}03$	$1.77\text{e-}02 \pm 6.45\text{e-}03$	0.178 ± 0.0092	8.90 ± 1.58
	
LFC 4400	0.50	$-1.40\text{e-}02 \pm 5.37\text{e-}03$	$-9.76\text{e-}03 \pm 5.11\text{e-}03$	0.422 ± 0.0090	0.09 ± 0.81
		$-3.47\text{e-}03 \pm 2.24\text{e-}03$	$-1.15\text{e-}02 \pm 2.09\text{e-}03$	0.435 ± 0.0036	4.24 ± 0.31
		$1.81\text{e-}02 \pm 6.89\text{e-}03$	$-1.20\text{e-}02 \pm 6.61\text{e-}03$	0.431 ± 0.0109	10.32 ± 0.95
		$-3.09\text{e-}02 \pm 1.44\text{e-}02$	$-1.21\text{e-}02 \pm 1.32\text{e-}02$	0.379 ± 0.0210	5.64 ± 1.97
LFC 4464	0.50
		$1.09\text{e-}02 \pm 1.33\text{e-}03$	$3.42\text{e-}03 \pm 1.24\text{e-}03$	0.119 ± 0.0030	15.46 ± 0.79
		$-5.17\text{e-}03 \pm 4.79\text{e-}03$	$3.00\text{e-}03 \pm 4.51\text{e-}03$	0.049 ± 0.0095	62.29 ± 4.46
		$4.89\text{e-}02 \pm 7.99\text{e-}03$	$2.50\text{e-}02 \pm 6.95\text{e-}03$	0.269 ± 0.0130	4.47 ± 1.61
LFC 4488	0.55
		$-5.27\text{e-}03 \pm 2.14\text{e-}03$	$1.77\text{e-}03 \pm 2.03\text{e-}03$	0.094 ± 0.0045	51.48 ± 1.48
		$3.12\text{e-}03 \pm 4.95\text{e-}03$	$2.07\text{e-}03 \pm 4.09\text{e-}03$	0.096 ± 0.0069	0.73 ± 2.16
	

Table 10—Continued

ID (1)	B/T (2)	$\langle \frac{a_3}{a} \rangle$ (3)	$\langle \frac{a_4}{a} \rangle$ (4)	$\langle \epsilon \rangle$ (5)	δPA (6)
LFC 4536	0.67	$2.43\text{e-}03 \pm 5.15\text{e-}03$	$-2.07\text{e-}02 \pm 4.04\text{e-}03$	0.015 ± 0.0116	0.37 ± 21.64
		$-5.58\text{e-}03 \pm 3.07\text{e-}03$	$-1.55\text{e-}02 \pm 2.28\text{e-}03$	0.058 ± 0.0065	28.22 ± 2.26
		$4.47\text{e-}02 \pm 1.55\text{e-}02$	$4.91\text{e-}02 \pm 1.31\text{e-}02$	0.237 ± 0.0146	0.68 ± 2.00
	
LFC 4654	0.92
		$1.90\text{e-}04 \pm 2.47\text{e-}03$	$-1.68\text{e-}02 \pm 1.89\text{e-}03$	0.035 ± 0.0054	28.63 ± 2.79
		$3.84\text{e-}02 \pm 9.79\text{e-}03$	$6.30\text{e-}02 \pm 9.47\text{e-}03$	0.228 ± 0.0123	5.13 ± 1.74
	
LFC 4657	0.89	$-8.52\text{e-}04 \pm 1.40\text{e-}03$	$-1.29\text{e-}03 \pm 1.16\text{e-}03$	0.024 ± 0.0032	10.61 ± 3.35
		$5.21\text{e-}03 \pm 1.67\text{e-}03$	$-6.97\text{e-}03 \pm 1.55\text{e-}03$	0.056 ± 0.0035	11.38 ± 1.83
		$-1.70\text{e-}02 \pm 6.71\text{e-}03$	$2.62\text{e-}02 \pm 6.02\text{e-}03$	0.152 ± 0.0122	4.19 ± 2.45
		$2.95\text{e-}03 \pm 2.49\text{e-}02$	$1.03\text{e-}01 \pm 3.71\text{e-}02$	0.063 ± 0.0453	16.61 ± 18.67
LFC 4681	0.92	$-9.28\text{e-}03 \pm 1.46\text{e-}03$	$1.57\text{e-}03 \pm 1.21\text{e-}03$	0.064 ± 0.0036	2.27 ± 1.69
		$-1.12\text{e-}02 \pm 7.62\text{e-}04$	$-2.53\text{e-}03 \pm 6.93\text{e-}04$	0.073 ± 0.0019	16.68 ± 0.76
		$-1.14\text{e-}02 \pm 2.15\text{e-}03$	$3.43\text{e-}03 \pm 2.09\text{e-}03$	0.109 ± 0.0045	5.52 ± 1.24
		$-3.45\text{e-}02 \pm 9.88\text{e-}03$	$-7.64\text{e-}03 \pm 8.92\text{e-}03$	0.123 ± 0.0175	3.75 ± 4.34
LFC 4740	0.74
		$-1.90\text{e-}02 \pm 8.92\text{e-}04$	$1.31\text{e-}03 \pm 7.81\text{e-}04$	0.113 ± 0.0028	12.13 ± 0.75
		$-9.20\text{e-}03 \pm 2.62\text{e-}03$	$4.71\text{e-}03 \pm 2.55\text{e-}03$	0.114 ± 0.0053	4.29 ± 1.40
		$-1.34\text{e-}02 \pm 6.86\text{e-}03$	$-2.53\text{e-}02 \pm 6.76\text{e-}03$	0.101 ± 0.0132	32.61 ± 3.87
LFC 4763	0.75
		$-1.49\text{e-}02 \pm 1.43\text{e-}03$	$1.42\text{e-}03 \pm 1.35\text{e-}03$	0.059 ± 0.0037	15.07 ± 1.74
		$-2.67\text{e-}02 \pm 4.29\text{e-}03$	$-1.90\text{e-}03 \pm 3.51\text{e-}03$	0.117 ± 0.0085	9.22 ± 2.04
		$-3.56\text{e-}02 \pm 1.08\text{e-}02$	$1.02\text{e-}02 \pm 9.23\text{e-}03$	0.163 ± 0.0186	39.44 ± 3.66

Note. — Column (1) gives our SExtractor catalog ID. Column (2) gives bulge-to-total luminosity ratio (see Section 5). Columns (3) to (6) give average isophotal parameters a_3/a , a_4/a , ellipticity and isophotal twist (in degree) in four different regions for each galaxy in four rows. Average value of isophotal parameter in Region 1, Region 2, Region 3 and Region 4 is listed in first, second, third and fourth row respectively for each galaxy.